

1187-114

NOV 13, 1974

1187-114

WHERE TO APPLY

Use in Food Handling and Processing Establishments • Stores • Dairies • Canneries • Hotels • Institutions • Hospitals • Schools • Homes

HOW TO APPLY

ROACHES, ANTS, SILVERFISH and CRICKETS:

Ascertain square feet of space to be treated. Open all doors, cabinets, closets, etc. Direct the mist toward insects, into cracks, crevices, and trails on door sills and window frames. For initial cleanup, apply for 10-15 seconds per 100 square feet of floor space. Close room for one hour. Repeat treatment as necessary. Do not spray directly on clothes, fabrics or other materials.

FLIES, MOSQUITOES, WASPS, GNATS: Ascertain cubic feet of space to be treated. Close doors and windows. Pointing nozzle upward, direct the aerosol mist to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion at least 36 inches from fabrics. Apply for 1 second per 1,000 cu. ft. of space. Spraying closer than 36 inches or longer than indicated may cause stains on fabrics, furs and other materials. Keep room closed 15 minutes after each treatment.

FLYING ADULT MOTHS: WEBBING CLOTHES MOTH, TOBACCO MOTH AND ALMOND MOTH — Ascertain cubic feet of space to be treated. Treat all suspected enclosures arranging clothing and other articles so that the foggy mist can circulate freely around them. Keep the nozzle in constant motion at least 36 inches away from the articles. Applying for 1 second per 1,000 cubic feet of space and close the treated area for 45 minutes or more. Spraying closer or longer than as directed above, or more often than bi-weekly, may cause stain on fabrics, furs and other materials. Two-Thirty-Two is not a moth proofing agent and will not destroy uncontacted adult moths.

STORED PRODUCT INSECTS (Accessible Stages):

(1) **CONFUSED FLOUR BEETLES, GRANARY WEEVIL, and RICE WEEVIL** — Direct treatment toward insects into cracks, crevices, around machinery, around containers of stored products and other areas that may harbor insects. Hit as many insects as possible. A minimum dosage of 10-15 seconds treating time per 100 sq. ft. of floor space is recommended, leaving treated area tightly closed for at least 45 minutes. Repeat treatment as required. (2) **INDIAN MEAL or CHOCOLATE MOTH, MEDITERRANEAN FLOUR MOTH and ANGIOLOUS GRAIN MOTH** — Close doors and windows. Direct aerosol to all parts of the room, especially into dark enclosures and around stored product containers, etc., where adult moths may be resting. Use a minimum dosage of 5 to 8 seconds treating time per 1,000 cu. ft. of space. Leave treated area closed 30 minutes. Repeat treatment weekly or as other flying moths appear.

Use in food processing plants only when plant is not in operation. All food processing surfaces should be covered or thoroughly cleaned before using. Food should be removed or protected during the treatment.

THIS CYLINDER IS
REFILLABLE

Return For Deposit When Empty

Made in U.S.A.

NET WT. 5 POUNDS
(2.268 KILOGRAMS)



Two Thirty Two

AEROSOL INSECTICIDE

KILLS ROACHES, ANTS, SILVERFISH, CRICKETS, FLIES, MOSQUITOES, WASPS, AND GNATS

**CAUTION: KEEP OUT OF
REACH OF CHILDREN**

See Side Panel for Additional Cautions

ACTIVE INGREDIENTS: Pyrethrins 1.95%, *Technical piperonyl butoxide 3.00%, N-Cetyl bicycloheptene dicarboximide 2.00%, Petroleum distillates 18.05%.

INERT INGREDIENTS: 75.0%

*Equivalent to 2.40% (butylcarbityl) (6-propylpiperonyl) ether and 0.60% of other related compounds.

WARNING: Contains under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator.

EPA Reg. No. 1187-114

VIRGINIA CHEMICALS INC., PORTSMOUTH, VA. 23703

How to assemble and operate the
VIRCHEM® Two-Thirty-Two
5 LB. CONTAINER — APPLICATOR

How to Assemble:

- (1) Remove valve protection cap from cylinder.
- (2) Insert cylinder into carrying bag.
- (3) Affix the hose and applicator to the valve with slight pressure from a wrench, if necessary.
- (4) Do not use without applicator.

How to Operate:

1. Sling strap of carrying bag over shoulder.
2. With applicator nozzle directed upward, open handwheel a full turn.
3. Keep the valve in the upright position at all times while in use.
4. Depress trigger on applicator handle to release spray.
5. Valves must be tightly closed and the cap replaced before shipment of empty cylinder.

CAUTION

- Do not spray directly on food or utensils.
- Do not remain in treated areas and ventilate the area after treatment is completed.
- Remove cats, birds and cover fish aquariums before spraying.
- Use in food processing plants only when plant is not in operation. All food processing surfaces should be covered or thoroughly cleaned before using. Food should be removed or protected.



LOT NO.