



POISON

**BEFORE USING THIS PRODUCT
READ THESE PRECAUTIONS**

- 1. LETHAL AIRE A-41** Insecticide CAN BE APPLIED ONLY WITH VIRGINIA CHEMICALS INC. AUTOMATIC EQUIPMENT. To insure proper dosage and safety, this product must be used with A-41 timer and applicator valves.
- 2. DO NOT CONTACT SKIN:** Natural rubber gloves impervious to DDVP must be worn. All skin surfaces must be covered with protective waterproof clothing before entering treated area or while changing aerosol units.
- 3. DO NOT INHALE DDVP MIST:** Wear a gas mask of a type passed by U.S. Bureau of Mines for DDVP protection if entering area during use period.
- 4. AEROSOL REPLACEMENT AND HANDLING RESTRICTED TO TRAINED PERSONNEL:** Due to the hazardous nature of this material only personnel trained by Virginia Chemicals Inc. should handle or replace it in service.
- 5. ADEQUATE PERSONAL HYGIENE AND CLEANLINESS** are necessary. After servicing remove clothing and bathe thoroughly with warm water and soap. Wash clothing before re-use.

ANTIDOTE AND FIRST AID TREATMENT

ATROpine is the emergency antidote for DDVP insecticide poisoning. Consult your physician about the need to have a supply of 1/2 grain atropine tablets for emergency use. **CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.**

INTERNAL: If the patient has been swallowed, induce vomiting immediately. This may be done by holding a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). **NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.**

EXTERNAL: If the material has been applied on the skin, immediately remove patient from the vicinity of the treated area and wash off all traces of the material. Then wash skin with soap and running water for at least 15 minutes.

IF WARNING SYMPTOMS APPEAR: Administer one 1/2 grain atropine tablet orally daily. **NEVER ADMINISTER ATROpine UNLESS WARNING SYMPTOMS APPEAR.** (See **WARNING SYMPTOMS** below).

- Call a physician immediately.
- Do not give the patient anything by mouth except water.
- Do not let the patient drink alcohol.

IMPORTANT -- SEE NOTE TO PHYSICIAN



POISON

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams until complete atropinization has occurred. 2-PAM is also antidotal and may be administered in conjunction with atropine.

- Morphine is contraindicated.
- Clear chest by postural drainage.
- **Observe patient continuously for 48 hours.**
- Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor.
- Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

The mechanical equipment used in the application of this insecticidal composition is covered by United States Patent Number 3,209,950, owned by Virginia Chemicals Inc.