

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.  
In case of spillage, do not use water for treatment or disposal.  
Do not use if secondary container is not marked with manufacturer's name and product name.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER**

**DO NOT CONTACT THE SKIN.** Gloves in percutaneous DDVP should be worn. All items of clothing should be immediately changed with protective clothing during application.  
**DO NOT INHALE DDVP MIST.** Wear a pesticide respirator (not a pump sprayer) approved by the Mine Safety and Health Administration and by the National Institute for Occupational Safety and Health for DDVP Protection.  
**ADEQUATE PERSONAL HYGIENE AND CLEANLINESS** are necessary. After applying remove clothing and wash thoroughly with warm water and soap. Wash clothing before reuse.  
**DO NOT CONTAMINATE FOOD.** Do not smoke, eat or chew tobacco in the operation area.

**FIRST AID TREATMENT**

**If swallowed:** Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit is clear. Do not induce vomiting if unconscious or if patient is unconscious.  
**If inhaled:** Remove victim to fresh air. Apply respiration if indicated.  
**If on skin:** Remove contaminated clothing and wash affected areas with soap and water.  
**If in eyes:** Flush eyes for at least 15 minutes with water. Call a physician immediately.  
**CALL A PHYSICIAN OR POISON CONTROL CENTER IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.** • Keep patient prone and quiet. • Start artificial respiration immediately if patient is not breathing. • Transport the patient immediately to the nearest physician.

**NOTE TO PHYSICIAN**

**WARNING SYMPTOMS:** Symptoms include weakness, headache, tightness in chest, blurred vision, runny nose, drooping eyelids, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.  
**TREATMENT:** Atropine is the specific therapeutic antagonist of the action of parasympathetic stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 1 to 2 minute intervals in doses of 1 to 2 mg. (grams) until complete atropinization has occurred.  
• Morphine is contraindicated. • Clear chest by postural drainage. • Observe patient continuously for 48 hours. • Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. • Allow a further 72 hour time for cholinesterase regeneration has been allowed (as determined by blood test).

ACCEPTED  
with COMMENTS  
in EPA Record Book

DEC 28 1982

1187-115