

ACCEPTED
2/15/78
FOR THE
FEDERAL AND
ECONOMIC
COMMISSIONS

FREEN

MINT ODORED DISINFECTANT

DISINFECTANT — CLEANER — SANITIZER —
FUNGICIDE — DEODORANT

ACTIVE INGREDIENTS:

N-Alkyl (50% C₁₂, 40% C₁₄, 10% C₁₆) dimethyl benzyl
ammonium chloride 4.8%
Sodium carbonate 3.5%
Essential Oils 0.3%

INERT INGREDIENTS: 91.4%

EPA Reg. No. 1190-43

DANGER:

KEEP OUT OF REACH OF CHILDREN

See First Aid and Other Precautions on Right Panel
Container Disposal: When empty, rinse container thoroughly with water and
destroy it by crushing or burying in a safe place.



USE DIRECTIONS

FOR CLEANING—First remove gross filth and heavy soil deposits.

Lightly soiled areas—surfaces to be cleaned should be mopped or brushed with a solution containing 1 ounce per gallon of water. Rinse cleaned area with fresh water.

Heavily soiled areas—mop or brush soiled surfaces with a solution containing 2-3 ounces per gallon of water. Rinse cleaned surface with fresh water.

GENERAL PURPOSE DISINFECTION—To disinfect walls, floors and other hard surfaces in homes, schools, restaurants, rest rooms and other public areas, mop or brush surface to be treated with a solution containing 1½ ounces per gallon of water. Surfaces that are heavily soiled should be cleaned with 2 ounces per gallon of water and then disinfected. Those surfaces which may later come in contact with food must be rinsed with clear water of drinking quality.

To disinfect locker rooms, shower stalls, walkways around pools and other areas where fungicidal action is required, mop area with a solution containing 2 ounces in 3 gallons of water. This solution will destroy *Trichophyton interdigitale*, a causative agent of athlete's foot.

HOSPITAL DISINFECTION—To disinfect walls, floors, furniture and other hard surfaces in hospitals and nursing homes, mop or brush surface to be cleaned with a solution containing 2½ ounces per gallon of water. Those surfaces that are heavily soiled with blood, fecal material or exudates should be thoroughly scrubbed with a solution containing 2 ounces per gallon of water and then rinsed with a fresh solution containing 2 ounces per gallon of water before disinfection.

MINT ODORED DISINFECTANT

DISINFECTANT — CLEANER — SANITIZER
FUNGICIDE — DEODORANT

ACTIVE INGREDIENTS

INERT INGREDIENTS

DANGER:

KEEP OUT OF REACH OF CHILDREN

See First Aid and Other Precautions on Right Panel

Container Disposal: When empty, rinse container thoroughly with water and destroy it by crushing or burying in a safe place.

SANITIZATION—To sanitize walls, floors and equipment in food processing plants, first clean thoroughly with a solution containing 1 ounce per gallon of water. If surface is heavily soiled, use at a rate of 2-3 ounces per gallon to insure thorough cleaning. Sanitize cleaned surface with a rinse solution containing ½ ounce per gallon of water. No rinse required after sanitization.

DEODORIZATION—To deodorize surfaces during cleaning, mop or brush surfaces with a solution containing 2 ounces per gallon of water.

This product is an effective anti-bacterial cleaner designed for use in household, industrial and institutional cleaning. When used as directed, it will disinfect hard surfaces, such as floors, walls, counter tops and furniture. It may be relied upon to deodorize bathroom areas, garbage pails, food storage areas, kennel areas and other areas where obnoxious odors may develop. This product may be used as a general purpose anti-bacterial — antifungal detergent in homes, schools, restaurants, hospitals, nursing homes, public rooms, food processing plants and in other places where efficient cleaning and antimicrobial action is desired.

DANGER—KEEP OUT OF REACH OF CHILDREN—Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID—In case of eye contact, immediately flush eyes with plenty of water for at least 15 minutes, then get medical attention promptly. In case of skin contact, flush skin with plenty of water for at least 15 minutes. Remove and wash contaminated clothing before reuse. If swallowed, drink immediately a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.