

10% MALATHION DUST

ACCEPTED
July 18, 1971
1191-175
With Comments

Active Ingredients:

Malathion (O,O-Dimethyl Dithiophosphate of Diethyl Mercaptosuccinate) - - - - - 10%

Inert Ingredients: - - - - - 90%

100%

ANTIDOTE

INTERNAL: ANTIDOTE AND FIRST AID: If any of the insecticide is swallowed, give tablespoonful of salt in glass of warm water and repeat until vomit fluid is clear. Take one or two 1/100 grain atropine tablets, and repeat in one hour if necessary. **CALL A PHYSICIAN IMMEDIATELY!** If splashed in eyes, wash immediately with water and call physician if irritation persists.

NOTE TO PHYSICIAN: In case of poisoning, atropine is antidotal. Observe patient carefully since repeated treatment may be needed. Symptoms are: Cholinesterase inhibition, headache, lacrimation, salivation, labored breathing, vomiting, marked tremors, diarrhea, convulsions. 2-PAM is also antidotal and may be administered in conjunction with atropine.

EXTERNAL: In case of skin contact, wash immediately with soap and water.

WARNING

KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed. Avoid inhaling or getting in eyes. Avoid repeated or prolonged skin contact. Wash thoroughly with soap and water after using. Keep clothing free from residues. Avoid contamination of feed and foodstuffs. Do not store near any product which will absorb odors. Keep out of reach of children and domestic animals. Malathion upon repeated or prolonged use may cause Cholinesterase inhibition.

Highly toxic to fish. Do not contaminate any body of water with direct application or washings from spray tank. Do not apply when weather conditions favor drift from area treated. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is for use only on cotton. Cotton crops which have received late applications of this insecticide should not be grazed by dairy animals or animals being finished for slaughter. Do not make applications within 7 days of picking cotton.

DIRECTIONS FOR USE

COTTON PESTS: To control aphids, spider mites, thrips dust at the rate of 10-15 lbs. per acre.

As local conditions may vary, consult your state agricultural experiment station or state extension service specialist for timing of applications and repeat as necessary.

USDA REG. NO. 1191 - 195

ACCEPTED WITH COMMENTS