

# FLIGHT BRAND



## NOTICE

The manufacturer guarantees and warrants that the active ingredient and the minimum active ingredient content and the net weight of the contents are as stated and that the directions, warnings and other statements on this label are reasonably and fairly based upon responsible testing of effectiveness, of toxicity to plants and to laboratory animals and of residue remaining on food crops. The manufacturer makes no other express or implied guarantee, warranty or representation whatever, of a similar or different sort and none is to be inferred by the purchaser or user. Neither Carolina Chemicals, Inc. nor the seller shall be held responsible in any manner for any personal injury or property damage or other type of loss resulting from the careless or improper handling, storage, or use of this material whether used singly or in combination with other materials. The buyer (or user) assumes and agrees to accept all risk and liability resulting from such handling, storage or use. No modification of this paragraph is authorized except by specific reference to this paragraph in writing by an officer of the manufacturer.

# FLIGHT

## BRAND DIRECTIONS FOR USE

Plant	Pest	Dust Per Acre	Residue Tolerance	Interval (days) Between Application and Harvest
Walnuts	Aphids	30-50 lbs.	8ppm	-0-
Blueberries	Blueberry Maggot	20 lbs.	8ppm	-1-
Cranberries	Leafhoppers	30 lbs.	8ppm	-3-
	Black-headed Fireworms	30 lbs.	8ppm	-3-
Grapes	Leafhoppers	16-32 lbs.	8ppm	-3-
	Spider mite	16-32 lbs.	8ppm	-3-
	Make 2 or more applications as needed.			
Strawberries	Aphids	40 lbs.	8ppm	-3-
	Spider mites	40 lbs.	8ppm	-3-
Beans	Mexican Bean Beetle	30-35 lbs.	8ppm	-1-
	Leafhopper	30-35 lbs.	8ppm	-1-
	Make 2 or more applications as needed.			
Broccoli	Aphids			-3-
Cabbage	Imported Cabbage-Worm	30 lbs.	8ppm	-7-
Kale	Cabbage Looper			-7-
Tuclid				-7-
Mustard Greens				-7-
Brussels Sprouts	Aphids	30 lbs.	8ppm	-7-
Carrots				
Radishes	Aphids	24-30 lbs.	8ppm	-7-
Parsnips				
Salsify				
Horseradish				
Cauliflower	Aphids	40 lbs.	8ppm	-7-
Collards				
Dandelion	Aphids	24-30 lbs.	8ppm	7
Watercress				
Swiss Chard				
Parsley				
Cucumbers	Aphids	30-35 lbs.	8ppm	-1-
Squash	Spider mites			
Melon:	Leafhopper on melon			
	Do not apply Malathion to Cucurbits unless plants are dry.			
Lettuce	Aphids	24-32 lbs.	8ppm	Leaf 14
	Mites			Head 7
	Cabbage Looper			
Onions	Thrips	30-40 lbs.	8ppm	-3-
Peas	Pea Aphid	25 lbs.	8ppm	-3-
Potatoes	Aphids	25-30 lbs.	8ppm	-3-
	Leafhopper			
Sprouts	Aphids	24-30 lbs.	8ppm	-7-
Tomatoes	Spider mites	35 lbs.	8ppm	-1-
	Aphids			
	Tomato Russet Mite			

**ORNAMENTALS:** Apply thoroughly as necessary to control Aphids, Spider mites, and Bagworms.

**POULTRY:** Dust individual birds by means of a shaker can or rotary hand duster to control Northern Fowl Mite and Poultry Lice. Repeat in 4-8 weeks or when necessary. As a litter treatment use 1 lb. per 50 sq. ft. of litter and floor space and to nests, roosts, and adjacent areas. Repeat as necessary. Avoid contamination of feed and feeding troughs and water fountains. Residue tolerance: Poultry - 4 ppm. Eggs - Zero ppm.


**DOGS & CATS:** Apply sufficient dust to treat animal thoroughly. Use any garden-type duster, or a special user-type dispenser. If the animal has free run, treatments may have to be made every second or third week. To avoid unnecessary applications, animal should be closely examined for fleas before retreating. If confined to an area free of fleas one treatment may be sufficient. For control of fleas on kennel animals, use 1 lb. per 100 sq. ft. and remove manure or debris. Dust area at rate of 1-2 lbs. per 1000 sq. ft. with rotary duster.

771

NET WT. 4 LBS.

# FLIGHT

BRAND




## ANTIDOTE

**INTERNAL: ANTIDOTE AND FIRST AID:** If any of the insecticide is swallowed, give tablespoonful of salt in glass of warm water and repeat until vomit fluid is clear. Take one or two 1/100 grain atropine tablets, and repeat in one hour if necessary. **CALL A PHYSICIAN IMMEDIATELY!** If splashed in eyes, wash immediately with water and call physician if irritation persists. **NOTE TO PHYSICIAN:** In case of poisoning, Atropine is antidotal. Observe patient carefully since repeated treatment may be needed. Symptoms are: Cholinesterase inhibition, headache, lacrimation, salivation, labored breathing, vomiting, marked tremors, diarrhea, convulsions. 2-PAAM is also antidotal and may be administered in connection with atropine.

**EXTERNAL:** In case of skin contact, wash immediately with soap and water.

# FLIGHT

BRAND



## 5% MALATHION DUST

**ACTIVE INGREDIENTS:** Malathion (O,O-Dimethyl Dithiophosphate of Diethyl Mercaptosuccinate)

**INERT INGREDIENTS:** .....

## WARNING

### KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed. Avoid inhaling or getting in eyes. Wash thoroughly with soap and water after using. Keep children and domestic animals out of treated areas. Avoid contamination of feed and foodstuffs. Do not store in areas where children and domestic animals will absorb odors. Keep out of reach of children and domestic animals. Do not use where milk is processed or stored. Do not apply in or near dairy barns. Remove animals and poultry from building before use. Re-use container. Destroy when empty.

Highly toxic to fish. Do not contaminate any body of water or washings from spray tank. Do not apply when weather is likely to cause runoff. Do not apply where runoff is likely to occur.

This product is highly toxic to bees exposed to direct contact with treated surfaces. Protective information may be obtained from your local Extension Service.

SEE SIDE PANEL FOR ANTIDOTE STATEMENT  
USDA REG. NO. 1191-192

**CAROLINA CHEMICALS, INC.**  
WEST COLUMBIA, S.C.

# FLIGHT BRAND

# FLIGHT BRAND®



## NOTICE

The manufacturer guarantees and warrants that the active ingredient and the minimum active ingredient content and the net weight of the contents are as stated and that the directions, warnings and other statements on this label are reasonably and fairly based upon responsible testing of effectiveness, of toxicity to plants and to laboratory animals and of residue remaining on food crops. The manufacturer makes no other express or implied guarantee, warranty or representation whatever, of a similar or different sort and none is to be inferred by the purchaser or user. Neither Carolina Chemicals, Inc. nor the seller shall be held responsible in any manner for any personal injury or property damage or other type of loss resulting from the careless or improper handling, storage, or use of this material whether used singly or in combination with other materials. The buyer (or user) assumes and agrees to accept all risk and liability resulting from such handling, storage or use. No modification of this paragraph is authorized except by specific reference to this paragraph in writing by an officer of the manufacturer.

# FLIGHT

BRAND®

## DIRECTIONS FOR USE

### VEGETABLES:


Plant	Pest	Dust Per Acre	Residue Tolerance	Interval (days) Between Last Application and Harvest
Walnuts	Aphids	30-50 lbs.	8ppm	-0-
Blueberries	Blueberry Maggot	20 lbs.	8ppm	-1-
Cranberries	Leafhoppers	30 lbs.	8ppm	-3-
	Black-headed fireworms	30 lbs.	8ppm	-3-
Grapes	Leafhoppers	16-32 lbs.	8ppm	-3-
	Spider mites	16-32 lbs.	8ppm	-3-
Make 2 or more applications as needed.				
Strawberries	Aphids	40 lbs.	8ppm	-3-
	Spider mites	40 lbs.	8ppm	-3-
Beans	Mexican Bean Beetle	30-35 lbs.	8ppm	-1-
	Leafhopper	30-35 lbs.	8ppm	-1-
Make 2 or more applications as needed.				
Broccoli	Aphids			-3-
Cabbage	Imported Cabbage-Worm	30 lbs.	8ppm	-7-
Kale	Cabbage Looper			-7-
Turnip				-7-
Mustard Greens				-7-
Brussels Sprouts	Aphids	30 lbs.	8ppm	-7-
Carrots				
Radishes				
Parsnips	Aphids	24-30 lbs.	8ppm	-7-
Salsify				
Horse radish				
Cauliflower	Aphids	40 lbs.	8ppm	-7-
Collards				
Dandelion				
Watercress	Aphids	24-30 lbs.	8ppm	-7-
Swiss Chard				
Parsley				
Cucumbers	Aphids	30-35 lbs.	8ppm	-1-
Squash	Spider mites			
Melons	Leafhopper on melon			
Do not apply Malathion to Cucurbits unless plants are dry.				
Lettuce	Aphids	24-32 lbs.	8ppm	Leaf 14
	Mites			Head 7
	Cabbage Looper			
Onions	Thrips	30-40 lbs.	8ppm	-3-
Peas	Pea Aphid	25 lbs.	8ppm	-3-
Potatoes	Aphids	25-30 lbs.	8ppm	-3-
	Leafhopper			
Spinach	Aphids	24-30 lbs.	8ppm	-7-
Tomatoes	Spider mites	35 lbs.	8ppm	-1-
	Aphids			
	Tomato Russet Mite			

**ORNAMENTALS:** Apply thoroughly as necessary to control Aphids, Spider mites, and Bagworms.

**POULTRY:** Dust individual birds by means of a shaker can or rotary hand duster to control Northern Fowl Mite and Poultry lice. Repeat in 4-8 weeks or when necessary. As a litter treatment use 1 lb. per 50-60 sq. ft. to litter and floor space and to nests, roosts and adjacent areas. Repeat as necessary. Avoid contamination of feed and feeding troughs and water fountains. Residue tolerance: Poultry—4ppm. Eggs—Zero ppm.

**DOGS & CATS:** Apply sufficient dust to treat animal thoroughly. Use any garden-type duster, puff duster or shaker-type dispenser. If the animal has free run, treatments may have to be made every second or third week. To avoid unnecessary applications, animal should be closely examined for fleas before retreating. If confined to an area free of fleas one treatment may be sufficient. For control of fleas in kennels, animal quarters, etc., remove manure or debris. Dust area at rate of 1-2 lbs. per 1,000 sq. ft. with shaker duster.

# FLIGHT BRAND®



## ANTIDOTE

**INTERNAL: ANTIDOTE AND FIRST AID:** If any of the insecticide is swallowed, give tablespoonful of salt in glass of warm water and repeat until vomit fluid is clear. Take one or two 1/100 grain atropine tablets, and repeat in one hour if necessary. **CALL A PHYSICIAN IMMEDIATELY!** If splashed in eyes, wash immediately with water and call physician if irritation persists. **NOTE TO PHYSICIAN:** In case of poisoning, Atropine is antidotal. Observe patient carefully since repeated treatment may be needed. Symptoms are: Cholinesterase inhibition, headache, lacrimation, salivation, labored breathing, vomiting, marked tremors, diarrhea, convulsions. 2-PAM also antidotal and may be administered in connection with atropine. **EXTERNAL:** In case of skin contact, wash immediately with soap and water.

# FLIGHT

BRAND

DIRECTIONS FOR USE

Pest	Dust Per Acre	Residue Tolerance	Interval (days) Between Last Application and Harvest
Aphids	30-50 lbs.	8ppm	-0-
Blueberry Maggot	20 lbs.	8ppm	-1-
Leafhoppers	30 lbs.	8ppm	-3-
Black-headed fireworms	30 lbs.	8ppm	-3-
Leafhoppers	16-32 lbs.	8ppm	-3-
Spider mites	16-32 lbs.	8ppm	-3-
Make 2 or more applications as needed.			
Aphids	40 lbs.	8ppm	-3-
Spider mites	40 lbs.	8ppm	-3-
Mexican Bean Beetle	30-35 lbs.	8ppm	-1-
Leafhopper	30-35 lbs.	8ppm	-1-
Make 2 or more applications as needed.			
Aphids			-3-
Imported Cabbage-Worm	30 lbs.	8ppm	-7-
Cabbage Looper			-7-
			-7-
			-7-
Aphids	30 lbs.	8ppm	-7-
Aphids	24-30 lbs.	8ppm	-7-
Aphids	40 lbs.	8ppm	-7-
Aphids	24-30 lbs.	8ppm	-7-
Aphids	30-35 lbs.	8ppm	-1-
Spider mites			
Leafhopper on melon			
Do not apply Malathion to Cucurbits unless plants are dry.			
Aphids	24-32 lbs.	8ppm	Leaf 14
Mites			Head 7
Cabbage Looper			
Thrips	30-40 lbs.	8ppm	-3-
Pea Aphid	25 lbs.	8ppm	-3-
Aphids	25-30 lbs.	8ppm	-3-
Leafhopper			
Aphids	24-30 lbs.	8ppm	-7-
Spider mites	35 lbs.	8ppm	-1-
Aphids			
Tomato Russet Mite			

Apply thoroughly as necessary to control Aphids, Spider mites, and Bagworms. Use individual birds by means of a shaker can or rotary hand duster to control Mites and Poultry lice. Repeat in 4-8 weeks or when necessary. As a litter treatment 0-60 sq. ft. to litter and floor space and to nests, roosts and adjacent areas. Repeat. Avoid contamination of feed and feeding troughs and water fountains. Residue 100-400 ppm. Eggs—Zero ppm.

Apply sufficient dust to treat animal thoroughly. Use any garden-type duster or sprayer-type dispenser. If the animal has free run, treatments may have to be repeated every second or third week. To avoid unnecessary applications, animal should be checked for fleas before retreating. If confined to an area free of fleas one treatment will suffice. For control of fleas on kennels, pens, stables, etc., remove manure or debris and apply 1-2 lbs. per 1,000 sq. ft. with hand duster.

NET WT. 4 LBS. AVDP.

# FLIGHT

BRAND



# FLIGHT

BRAND



### ANTIDOTE

**INTERNAL: ANTIDOTE AND FIRST AID:** If any of the insecticide is swallowed, give tablespoonful of salt in glass of warm water and repeat until vomit fluid is clear. Take one or two 1/100 grain atropine tablets, and repeat in one hour if necessary. **CALL A PHYSICIAN IMMEDIATELY!** If splashed in eyes, wash immediately with water and call physician if irritation persists. **NOTE TO PHYSICIAN:** In case of poisoning, Atropine is antidotal. Observe patient carefully since repeated treatment may be needed. Symptoms are: Cholinesterase inhibition, headache, lacrimation, salivation, labored breathing, vomiting, marked tremors, diarrhea, convulsions. 2-PAM is also antidotal and may be administered in connection with atropine.

**EXTERNAL:** In case of skin contact, wash immediately with soap and water.

## 5% MALATHION DUST

ACTIVE INGREDIENTS:  
 Malathion (O,O-Dimethyl Dithiophosphate of Diethyl Mercaptosuccinate) ..... 5%  
 INERT INGREDIENTS: ..... 95%  
 100%



### WARNING

#### KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed. Avoid inhaling or getting in eyes. Avoid skin contact. Wash thoroughly with soap and water after using. Keep clothing free from residues. Avoid contamination of feed and foodstuffs. Do not store near any product which will absorb odors. Keep out of reach of children and domestic animals. Malathion upon repeated or prolonged use may cause Cholinesterase inhibition. Do not use where milk is processed or stored. Do not apply on dairy animals or in dairy barns. Remove animals and poultry from buildings before treating. Do not re-use container. Destroy when empty.

Highly toxic to fish. Do not contaminate any body of water with direct application or washings from spray tank. Do not apply when weather conditions favor drift from area treated. Do not apply where runoff is likely to occur.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

SEE SIDE PANEL FOR ANTIDOTE STATEMENT  
 USDA REG. NO. 1191-198

**CAROLINA CHEMICALS INC.**  
 WEST COLUMBIA, S.C.