

DIRECTIONS FOR USE

ORNAMENTALS — Thorough, full-coverage applications should be made. Repeat applications as necessary:

TEASPOONFUL PER GALLON	TEASPOONFUL PER GALLON
Aphids	1½
Azalea scale	2
Bagworms	2
Birch leaf miner	2
Boxwood leaf miner	2
Euonymus scale	1½
European pine shoot moth	1½
Fletcher scale*	2
Florida red scale*	2
Four-lined leaf bug	1½
Japanese beetle (adult)	1½
Juniper scale*	2
Lace bug	1
Magnolia scale	2
Mealybugs	1½
Monterey pine scale	2½
Oak Kermes*	2
Oyster shell scale*	1
Pine leaf scale	2
Pine needle scale*	4
Potato leaf hopper	1½
Rose leaf hopper	1½
Scurfy scale	1½
Soft scale	2½
Spider mites	1½
Tarnished plant bug	1½
Tent caterpillar	2
Thrips	1½
Whitefly	1½
Black scale crawlers	2¼

*Apply when scale crawlers have settled on foliage. At 4 teaspoons/gal. rate, injury may occur on certain ferns and on some species of crassula and canaerti juniper.

For millipedes springtails and sowbugs, use 1 teaspoonful per gallon of spray and apply to 150 sq. ft. of soil surface or where insects congregate.

LAWNS — Against ground pearls use 1 teaspoonful per gallon of spray per 5,500/7,000 sq. ft. Make full-coverage application to soil surface when ground pearl nymphs are in the pink, "crawler" or active stage and immediately wash into soil with additional water.

TOBACCO — For control of Aphids (plant lice) on tobacco thoroughly mix 1 pint of this material to 60 gallons of water and spray the resulting mixture at the rate of 40 to 80 gallons per acre. Be sure to obtain good coverage over the entire plant. Do not apply to tobacco within 7 days of harvest.

FLEAS ON DOGS AND CATS — Use 2 oz. per gallon of water. Wet animals thoroughly using a hand sprayer. Repeat treatments every 2 or 3 weeks, if necessary. **DO NOT TREAT ANIMALS UNDER ONE MONTH OF AGE.** In **ANIMAL QUARTERS** use 5 oz. per gallon of water and use 1 gallon of resulting spray per 1,000 sq. ft. Use garden sprayer. Remove debris, then treat and repeat when necessary.

1 GALLON

FLIGHT
BRAND®



57% MALATHION

(Contains 5 lbs. Malathion p

ACTIVE INGREDIENTS:

Malathion

Xylene

INERT INGREDIENTS:

WARNING: KEEP OUT OF REAC

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONS

Applications as necessary:

TEASPOONSFUL PER GALLON
..... 2½
..... 2
..... 1
..... 2
..... 4
..... 1½
..... 1½
..... 1½
..... 2½
..... 1½
..... 1½
..... 2
..... 1½
..... 1½
..... 2½

..... on some species of

..... soil surface or where insects

..... coverage application to soil
..... soil with additional water.

..... gallons of water and spray
..... entire plant. Do not apply

..... Repeat treatments every
..... ERS use 5 oz. per gallon of
..... and repeat when necessary.

1 GALLON

FLIGHT

BRAND®



57% MALATHION EC

(Contains 5 lbs. Malathion per gallon)

ACTIVE INGREDIENTS:

Malathion	57%
Xylene	38%
INERT INGREDIENTS:	5%
	<u>100%</u>

WARNING: KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONS & ANTIDOTE STATEMENT.

ACCEPTED
5/28/71
FOR SALE IN THE DISTRICT OF COLUMBIA
BY LICENSE NO. 1191-226

HARMFUL IF SWALLOWED. FLAMMABLE. Avoid breathing. Keep children off treated areas.

Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of lakes, streams or contaminate water by cleansing of equipment, or disposal

Do not reuse empty container. Wash empty container th

ANTIDOTE: If swallowed, induce vomiting with warm sal immediately wash skin with soap and water.

CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPE appetite, nausea, vomiting, abdominal cramps, diarrhea, i restlessness, give two atropine tablets (1/100 gr. each) at c present. Have patient lie down and keep quiet.

PHYSICIAN'S NOTE: Warning symptoms include loss of sweating, constriction of pupils, blurred vision, giddiness,

TREATMENT: Give 1/30 to 1/40 gr. atropine, preferably i every hour as required up to 30 tablets or until pupils dila respiration or oxygen administration may be necessary.

terase inhibitors may, without warning, cause prolonged st no further exposure until time for cholinesterase regenerat may be used in combination with atropine under controlle

NOTICE: The manufacturer guarantees and warrants that the active ingred stated and that the directions, warnings and other statements on this label c plants and to laboratory animals and of residue remaining on food crops. T whatever, of a similar or different sort and none is to be inferred by the i in any manner for any personal injury or property damage or other type of whether used singly or in combination with other materials. The buyer (or u or use. No modification of this paragraph is authorized except by specific

Manufactured By

CAROLINA
USDA REG



ACCEPTED
 5/28/71
 FEDERAL REGISTER
 REGISTRATION NO. 1191-226

HARMFUL IF SWALLOWED. FLAMMABLE. Avoid breathing spray mist. Avoid contact with skin. Wash thoroughly after using. Keep children off treated areas.
 Avoid contamination of feed and foodstuffs.
 This product is toxic to fish. Keep out of lakes, streams or ponds. Apply this product only as specified on this label. Do not contaminate water by cleansing of equipment, or disposal of wastes.
 Do not reuse empty container. Wash empty container thoroughly with water and detergent. Discard in safe place.

ANTIDOTE: If swallowed, induce vomiting with warm salty or soapy water. If on the skin, remove contaminated clothing and immediately wash skin with soap and water.

CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING: If symptoms or signs of poisoning include loss of appetite, nausea, vomiting, abdominal cramps, diarrhea, excessive sweating, constriction of pupils, blurred vision, giddiness, and restlessness, give two atropine tablets (1/100 gr. each) at once. Remove patient immediately from the area where Malathion is present. Have patient lie down and keep quiet.

PHYSICIAN'S NOTE: Warning symptoms include loss of appetite, nausea, vomiting, abdominal cramps, diarrhea, excessive sweating, constriction of pupils, blurred vision, giddiness, and restlessness.

TREATMENT: Give 1/30 to 1/40 gr. atropine, preferably in injection, or two or three tablets at once and parenterally or orally every hour as required up to 30 tablets or until pupils dilate. Never give morphine. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood tests. NOTE: 2-PAM may be used in combination with atropine under controlled conditions. Refer to 2-PAM directions.

NOTICE: The manufacturer guarantees and warrants that the active ingredient and the minimum active ingredient content and the net weight of the contents are as stated and that the directions, warnings and other statements on this label are reasonably and fairly based upon responsible testing of effectiveness, of toxicity to plants and to laboratory animals and of residue remaining on food crops. The manufacturer makes no other express or implied guarantee, warranty or representation whatever, of a similar or different sort and none is to be inferred by the purchaser or user. Neither Carolina Chemicals, Inc. nor the seller shall be held responsible in any manner for any personal injury or property damage or other type of loss resulting from the careless or improper handling, storage or use of this material whether used singly or in combination with other materials. The buyer (or user) assumes and agrees to accept all risk and liability resulting from such handling, storage or use. No modification of this paragraph is authorized except by specific reference to this paragraph in writing by an officer of the manufacturer.

NEC
 gallon)

57%
 38%
 5%
 100%

OF CHILDREN
 ANTIDOTE STATEMENT.

Manufactured By CAROLINA CHEMICALS, INC. West Columbia, S. C.
 USDA REG. NO. 1191-226