

# SEVIN®-PARATHION

## Tabacco Dust 10-1

### ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) -----


10% by wt.

Parathion (O, O-diethyl O-p-nitrophenyl thiophosphate) -----

1% by wt.

### INERT INGREDIENTS: -----

89% by wt.



**KEEP OUT OF REACH OF CHILDREN**

MAY BE FATAL IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN! RAPIDLY ABSORBED THROUGH SKIN! DO NOT GET IN EYES, ON SKIN OR ON CLOTHING. WEAR NATURAL RUBBER gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Dept. of Agriculture for parathion protection. Keep all equipment persons out of operating areas or vicinity where they may be danger of drift. Vacated areas should not be re-entered until drifting insecticides and volatile residues have dissipated. Workers entering a treated field within 7 days of an application should wear protective clothing. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and hot water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply when wind is blowing toward water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT FOR USE OR STORAGE IN OR AROUND THE HOME. DO NOT RE-USE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTES OR BURNING. IF BURNED STAY AWAY FROM SMOKE OR FLAME.

## DIRECTIONS FOR USE

READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS

For hornworms, budworms, aphids, flea beetles, suckfly, grasshoppers and Japanese beetle, apply 15 to 25 pounds per acre depending on the size of the plants. Make applications when insect first appears on the crop. Repeat applications as necessary. Avoid excessive applications.

For budworms, when dust applications are directed to the bud and top leaves only, apply 4 to 6 pounds per acre. Wear natural rubber gloves, protective clothing and goggles. Wear a mask or respirator of a type passed by the U. S. Dept. of Agriculture for Parathion protection while performing these operations. Do not apply with cloth bags.

Do not apply within five days of priming or fifteen days of cutting. Avoid getting wet with plant juices when cutting the crop.

®Trade Mark of UCC for 1-naphthyl N-methylcarbamate.