BLIGHT BRAND

DIRECTIONS FOR USE

Apply thoroughly at first sign of infestation and repeat as necessary

Plant	Pest	Dust Per Acre	Residue Tolerance	Days between last Application & Harvest
Beans	Mexican Bean Beetle	5 lbs.	10 ppm	0
	Leaf hoppers, bean leaf beetle, cucumber beetle, Western bean cutworm, flea beetles & Japanese beetle	10 lbs.	10 ppm	0
	Corn earworm, army worms, velvet bean caterpillar	10-15 lbs.	10 ppm	0
	Lima bean pod borer, Lygus and stink bugs	20 lbs.	10 ppm	0
Cucumbers Summer Squash	Pickle worm & melonworm	5 lbs.	10 ppm	0
	Cucumber beetles, squash bugs and flea beetles	10 lbs.	10 ppm	0
Sweet Corn	Corn earworm,* European Corn borer,* fall army worm, Sap beetles & Japanese beetle	10-20 lbs	10 ppm	Day of harvest for fresh market ears. 7 days before harvest on forage & husks fed to livestock
	*When attacking tassels, apply to [illegible] repeat as necessary. When attacking [illegible] apply liberally at 2 to [illegible] when the first silk appears [illegible] begin to dry. Three or more [illegible] may not depend [illegible] of the protection [illegible] during the pollination period [illegible] per application.			
Tomatoes Eggplant Peppers	Tomato fruitworm, fall army worm, flea beetles & leaf hoppers	10 lbs	10 ppm	0
	Tomato hornworm, European Corn borer & stink bugs	20 lbs	10 ppm	0
Potatoes	Colorado potato beetle	5 lbs	10 ppm	0
	Leaf hoppers & flea beetles	10 lbs	10 ppm	0
	Lygus bugs	20 lbs	10 ppm	0
	[illegible] ANC [illegible] at [illegible] 24 [illegible]	[illegible] to 2 [illegible]	ppm [illegible]	[illegible]

NOTE FOR PHYSICIAN:
CARBARYL IS A MODERATE, CHOLINESTERASE INHIBITOR. ATROPINE IS ANTIDOTAL.

S 71

ACCEPTED
MAR 21, 1973
[illegible]
TO ATTACHED COMMENTS