

COPPER-SULPHUR PEANUT

7.5 - 85

JAN 22 1971
1191-298

ACTIVE INGREDIENTS:

Copper (in the form of basic copper sulfate) expressed as Metallic - - - - -
Sulphur - - - - -

INERT INGREDIENTS - - - - -

Total - - - - -

CAUTION

KEEP OUT OF REACH OF CHILDREN: Harmful if swallowed. Avoid inhalation


DIRECTIONS FOR USE

For control of Peanut Leaf Spot or Cercospora Leaf Spot of Peanut.

1. Begin when the first spots appear on the basal leaves, usually 60-65 days after planting or June 15 to July 1.
2. Make 3 applications at intervals of 2 weeks, unless the first applications must be repeated because of rain, in which case 4 applications protection.
3. Apply 15 to 20 lbs. dust per acre, depending upon the size of the plant and the efficiency of the dusting equipment.

ER-SULPHUR PEANUT DUST

7.5 - 85



basic copper sulfate) expressed as Metallic	- - - - -	4.0%
- - - - -	- - - - -	85.0%
- - - - -	- - - - -	11.0%
- - - - -	- - - - -	<hr/> 100.0%

CAUTION

USDA REG. NO. 1191 - 278

KEEP OUT OF REACH OF CHILDREN: Harmful if swallowed. Avoid inhalation of dust.

DIRECTIONS FOR USE

or Cercospora Leaf Spot of Peanut.

ear on the basal leaves, usually 60-65 days after planting or June 15 to July 1.

als of 2 weeks, unless the first applications must be repeated because of rain, in which case 4 applications may be required for adequate

re, depending upon the size of the plant and the efficiency of the dusting equipment.