
DIRECTIONS FOR USE

MAY 3 0 1975

Under the Federal insection 18, Fungicide, and Areas nucles Act, as amended, for the preside registered under EPA Reg. No. /202

DIREGRO



BEANS: Lima bean pod borer, apply 50 lbs. per acre. For corn earworm, lygus and stink bugs, apply 38 lbs. per acre. Do not apply within 21 days of harvest.

PEPPERS: For control of tomato fruitworm, fall arrayworm, tomato hornworm, stink bug. Apply 50 lbs. per acre. Do not apply within 15 days of harvest.

TOMATOES: For control of tomato fruitworm, fall armyworm, tomato hornworm, stink bug. Apply 50 lbs. per acre. Do not apply within 15 days of harvest.

LETTUCE: For control of corn earworm, armyworm, lygus and stink bugs. Apply 50 lbs. per acre. Do not apply within 21 days of harvest. To avoid possible injury do not apply when foliage is wet or excessive humidity is expected during the next 2 days.

Empty bags should be buried or burned. (Stay away from smoke). If the bags are burned, the ashes should be buried in a safe place.

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)* **(O,O-dimethyl O-p-nitrophenylphosphorothioate INERT INGREDIENTS:

Copyright Union Carbide Corporation *U.S. Patent No. 2,903,478 **Methyl Parathion

E.P.A. Reg. No. 1202-298-AA

WARNING! **KEEP OUT OF REACH OF CHILDREN**

May be fatal if swallowed, inhaled, or absorbed through skin. Rapidly absorbed thru the skin. Do not get in eyes, on skin, or on clothing. Wear natural rubber gloves. protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for Sevin and Methyl Parathion protection. Keep all unprotected persons out of operating of drift. Vacated area should not be re-entered until drifting insecticides and volatile residues have dissipated.



NET WEIGHT SOLRS.

WARNING

SEVIN® METHYL -298 PARATHION 4-2 DUST

Do not contaminate feed and foodstuffs. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and water before re-use.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

.94.00%

100.00%

.4.00%

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

CONDITIONS OF SALE:

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.