




**TOXAPHENE METHYL
PARATHION 7E3E**



CONDITIONS OF SALE:

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, and handling of this material not in strict accordance with directions given herewith.




LOS ANGELES CALIFORNIA

NET CONTENT _____ GALS.

LOT NO.

475 CN



**FLIGHT BRAND
TOXAPHENE 80S**

55-gal net

An Insecticidal Toxicant

ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene containing 67-69% Chlorine) .. 80%

INERT INGREDIENTS: 20%
100%

FOR MANUFACTURING USE ONLY

WARNING

KEEP OUT OF REACH OF CHILDREN.

IN CASE OF FIRE, LEAK OR OTHER EMERGENCY - REPORT AT ONCE BY TELEPHONE TO:

CHEMTREK (AREA CODE 800/424-9300)

Because of the possibility of poisonous material adhering to the container, it must be thoroughly cleaned before use for materials other than poisons.

An Insecticide for Formulating Use

This material contains 80% Toxaphene by weight and 20% Xylene type solvent by weight in liquid state. Except in extremely low temperatures this material is freely flowable and may be used either in the formulation of liquid concentrates of Toxaphene and/or Toxaphene combined with other active ingredients (such as parathion, methyl parathion, malathion) or may be impregnated on an absorbant dust carrier to yield a finished dust or dust base concentrate.

Formulators using this product are responsible for providing data to support their label registrations.

Manufactured by
CAROLINA CHEMICALS, INC.
West Columbia, S. C. 29169

EPA Reg. No. 1191-340

EPA-EST. 1191-SC-1

FIRST AID - ANTIDOTE

Internal - Give Sodium Chloride 15g. (one tablespoonful of table salt) in a glass of warm water and repeat until vomit is clear,

External - Wash with soap & water.

Call a Physician Immediately

Physician's Note: Toxaphene produces central nervous system excitation which responds to barbituric acid derivatives. Prevent or control convulsions with barbiturates, preferable administered intravenously. If the barbiturate can be given before convulsions start, sodium pentobarbital is the drug of choice. Since toxaphene has a fairly long duration of action, a long-acting barbiturate such as phenobarbital may be used after initial control of convulsions. Refer to "Clinical Handbook on Economic Poisons," which was prepared by the U.S. Dept. of Health, Education and Welfare, Public Health Service, Communicable Disease Center, Atlanta, Georgia.

May be fatal if taken internally. Do not breathe vapors. Do not get on skin or in eyes or on clothing. Can be absorbed through skin in dry form or in solution. Keep clothing free from residue. Wash thoroughly after using. Avoid contamination of feed and food-stuffs. Do not store near food or feedstuffs.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

Not for use or storage in or around the home. Do not use, pour, spill or store near open flame.

Toxic to fish, wildlife and birds. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes.