

OCT 16 1967

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 24411 SUBJECT
~~TO BE REMOVED~~ COMMENTS

RECOMMENDATIONS

Apply when air is calm using power dusting equipment or aircraft. For good control make complete coverage of plant surfaces, treating both upper and lower leaf surfaces. Apply when mites first appear and repeat at 7-10 day intervals, or as needed to maintain control, but do not use closer to harvest than specified.

DIRECTIONS FOR USE

Fruit and Nuts Crops

APPLES, PEARS: To control two-spotted, Schoene, McDaniel, yellow (carpini), Willamette, Pacific and apple rust mites; for clean up of European red mite and clover mite--Apply 40-50 lbs. per acre at petal fall to second cover or whenever mite population appears. Do not apply within 7 days of harvest.

CHERRY, NECTARINES, PEACHES, PLUMS, PRUNES: European red, clover (brown almond), two-spotted, Schoene, McDaniel, Pacific, Willamette, peach silver, plum nursery and yellow mites--Apply 40-50 lbs. per acre at petal fall to first cover or whenever mite population appears. If repeat application is necessary, wait 30 days to avoid excessive residue. Do not apply within 7 days of harvest on cherries, plums and prunes; 14 days on peach and nectarine.

GRAPES: Brown almond (clover), European, McDaniel, Pacific, two-spotted, Willamette and yellow mites--Apply 20-35 lbs. per acre. Do not apply within 7 days of harvest.

RASBERRIES, DEWBERRIES, LOGANBERRIES, BLACKBERRIES, BOYSENBERRIES: Brown almond (clover), two-spotted, European red, Pacific, McDaniel, Willamette and yellow mites--Apply 15-30 lbs. per acre. Do not apply within 2 days of harvest.

STRAWBERRIES: Cyclamen, two-spotted, strawberry (Atlantic) and six-spotted mites--Apply 25-35 lbs. per acre. Do not apply within 2 days of harvest.

WALNUTS, FILBERTS: Brown almond (clover), two-spotted, European red, Pacific, McDaniel, Willamette and yellow mites--Apply 40-50 lbs. per acre. Do not apply within 14 days of harvest. Do not feed hulls to livestock.

Vegetable and Field Crops

ALFALFA, CLOVER AND OTHER LEGUMES FOR SEED: Two-spotted, strawberry (Atlantic), carmine and Pacific mites--Apply 25-35 lbs. per acre. Do not feed straw or threshings to dairy animals or animals being finished for slaughter.

Lot No. _____ R22367R1W



KELTHANE® SULFUR 4-50 DUST

AN AGRICULTURAL MITICIDE

ACTIVE INGREDIENTS:	
1, 1-bis (p-chlorophenyl)-2, 2, 2-Trichloroethanol	4.0%
Sulfur	50.0%
INERT INGREDIENTS:	46.0%
	100.0%

• Registered trademark of Rohm & Haas Company

CAUTION

KEEP OUT OF REACH OF CHILDREN.

Avoid repeated skin contact. Avoid inhalation of dust. Avoid contamination of feed and foodstuffs. Wash thoroughly after handling.

OBSERVE THE FOLLOWING USE LIMITATIONS

- Observe use limitations given on the label for specific crops.
- Do not use on feed or forage to be fed to dairy animals or animals being finished for slaughter.
- This product is toxic to fish. Keep out of any body of water.
- The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, only at recommended rates, and do not apply closer to harvest than specified.



Net Weight 50 lbs.

(Continued from left panel)

BEANS: Two-spotted, strawberry mites--Apply 20-35 lbs. per acre. Do not feed treated forage to dairy or meat animals within 7 days of harvest.

COTTON: Atlantic (strawberry) mites--Apply 20-35 lbs. per acre depending on size of plant. Do not feed treated forage to dairy or meat animals.

HOPS: Two-spotted mites--Apply 20-35 lbs. per acre. Do not feed treated forage to dairy or meat animals.

POTATOES: Two-spotted, European red mites--Apply 20-35 lbs. per acre.

SUGAR BEETS: Two-spotted, European red mites--Apply 20-35 lbs. per acre. Do not apply within 7 days of harvest to dairy or meat animals.

TOMATOES, EGGPLANT, PEPPERS: Two-spotted mites--Apply 20-35 lbs. per acre.

ORNAMENTALS, FLOWERS: European red, Pacific, Willamette and yellow mites in greenhouse or outdoors--Apply 20-35 lbs. per acre. Do not apply if mites are present and repeat as needed.

CONDITIONS OF SALE: 1. That this material conforms to the specifications and is reasonably fit for use as directed. 2. That this material is not authorized by any agent or representative of the manufacturer to be used for any purpose other than that of FITNESS or of MERCHANTABILITY, express or implied, concerning 2. Critical and unforeseeable risks which cannot be prevented by the elimination of chemicals. Such risks include damage to crops to which the material is applied, and damage caused by drift to other crops, though the product is reasonable in use. 3. Though label directions are followed, the user assumes all risks and liability (under 1 above) resulting from the use of this material.

USDA Reg. No. 1202-160

TURN EX
KEEP OUT

OCT 16 1967

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 1222-100 SUBJECT
~~TO PREPARE COMMENTS~~

(Continued from left panel)

BEANS: Two-spotted, strawberry (Atlantic), carmine and Pacific mites - Apply 20-30 lbs. per acre. Do not feed treated vines to meat or dairy animals. Do not apply within 7 days of harvest.

COTTON: Atlantic (strawberry), Pacific, two-spotted and desert mites--Apply 15-35 lbs. per acre depending on size of plants. Do not apply within 14 days of harvest or feed treated forage to dairy or meat animals.

HOPS: Two-spotted mites - Apply 20-30 lbs. per acre during early season; 30-35 lbs. for mid to late season. Do not apply within 7 days of harvest.

POTATOES: Two-spotted, Pacific and strawberry (Atlantic) mites - Apply 20-35 lbs. per acre.

SUGAR BEETS: Two-spotted, Pacific, strawberry (Atlantic) mites - Apply 15-30 lbs. per acre. Do not apply within 21 days of harvest. Do not feed crop residues (tops) to dairy or meat animals.

TOMATOES, EGGPLANT, PEPPERS: Pacific, tropical, tomato russet and two-spotted mites - Apply 20-35 lbs. per acre. Do not apply within 2 days of harvest.

ORNAMENTALS, FLOWERS, NURSERY STOCK AND SHADE TREES: For control of mites in greenhouse or outdoor plants, make full coverage applications when mites are present and repeat as necessary.

CONDITIONS OF SALE: 1. PureGro Company (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.
2. Critical and unforeseeable factors beyond the manufacturer's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above) resulting from handling, storage, and use of this material.

USDA Reg. No. 1202-100

TURN EMPTY CONTAINER
KEEP OUT OF SMOKE

4-50 DUST

ACICIDE

2-Trichloroethanol	4.0%
.....	50.0%
.....	46.0%
.....	100.0%

of Rohm & Haas Company

CHILDREN.

ation of dust. Avoid
sh thoroughly after handling.

E LIMITATIONS

l for specific crops.

dairy animals or animals being finished

l any body of water.

in has set official tolerances for
residues on food and forage crops
ly at recommended rates, and do