Page 1 of 1 Pg# 1202-107 PM 21

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER

Causes irreversible eye damage. May cause allergic skin reactions. Harmful if swallowed or inhaled. Do not get in eyes. Wear goggles or face shield and rubber gloves when handling. Avoid contact with skin and clothing. Avoid breathing of dust or vapor. Wash hands after use. Remove and launder clothing before reuse. Do not contaminate feed or foodstuffs.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

REENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.
Do not enter treated areas without protective clothing until dusts have settled.
Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.
When oral or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given where there is reason to believe that written warnings cannot be understood by workers. Written or oral warnings must include the following information:

"DANGER" "Area to be treated with captan on (date of application). Do not enter without appropriate protective clothing until dusts have settled. If accidentally exposed, consult the Statement of Practical Treatment portion of the pesticide label for first aid and medical attention recommendations."

PureGro
CAPTAN DUST 10

ACTIVE INGREDIENT:

*Captan (N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide)	10.0%
INERT INGREDIENTS:	90.0%
	100.0%

*U.S. Patent Numbers 2,553,770 and 2,553,771.

EPA REG. NO. 1202-107-AA
E.P.A. Est. No.

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

STATEMENT OF PRACTICAL TREATMENT

If in eyes—immediately flush eyes with plenty of water for at least 15 minutes. Call a physician.

If swallowed—drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious person. Get medical attention.

If inhaled—remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.

If on skin—wash skin with soap and water.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.
Open dumping is prohibited.

STORAGE

Keep pesticide in original container.
Store in cool, dry place.
Protect from excessive heat.
Do not store or transport near food or feed.
For help with any spill, leak, fire or exposure involving this material, call day or night (800) 424-9300.

PESTICIDE DISPOSAL

This product is acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Completely empty bag into application equipment. Do not reuse container. Dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

PUREGRO CONDITIONS OF SALE AND WARRANTY

PUREGRO COMPANY warrants only that this product conforms to the chemical description on the label and is reasonably fit for the uses stated on the label when used in strict accordance with the directions under normal conditions of use. Control and ineffectiveness failures because of PUREGRO COMPANY's control product have always arisen from causes beyond the control of PUREGRO COMPANY, including, but not limited to, storage or planting and weather which may affect the product or the application thereof. All such risks are assumed by the buyer. PUREGRO COMPANY shall not be liable for any loss or damage to plants and crops to which this product is applied, whether or not in accordance with the directions, unless such loss or damage is caused by gross negligence of PUREGRO COMPANY. PUREGRO COMPANY shall not be liable for any loss or damage resulting from the handling or use of this product unless PUREGRO COMPANY is notified of the claim within 8 days after the buyer discovers or should have discovered the loss or damage, and PUREGRO COMPANY shall not be liable for any consequential damages. PUREGRO COMPANY MAKES NO WARRANTY OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, EXPRESS OR IMPLIED, EXCEPT AS STATED ON THIS LABEL. Buyer and user assume the risk of any use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseen by PUREGRO COMPANY. Buyer and user acknowledge and assume all risks and liabilities resulting from the handling, storage and use of this product not in accordance with the directions, or under abnormal conditions, or under conditions not reasonably foreseen by PUREGRO COMPANY. PUREGRO COMPANY'S liability, whether in contract, in tort, under any warranty, in negligence or otherwise, shall not exceed the purchase price paid by the user for the product, and upon the return of the product, the purchase price paid by the user will be refunded. PUREGRO COMPANY SHALL NOT BE RESPONSIBLE FOR CONSEQUENTIAL DAMAGES. THE FOREGOING CONDITIONS OF SALE AND WARRANTY CAN BE VARIED ONLY BY AN AGREEMENT IN WRITING SIGNED BY A DULY AUTHORIZED REPRESENTATIVE OF PUREGRO COMPANY.

CROP RECOMMENDATIONS

Apply when air is calm with an efficient dust applicator, securing complete coverage with a light film of dust. Begin at first sign of infection and repeat at 5-10 day intervals as needed to maintain control. Under severe disease conditions use the higher rate.

FRUIT AND NUT CROPS

ALMONDS Apply 40-50 lbs per acre to control Brown Rot, Shot Hole, Scab, and Leaf Blight. Apply at popcorn, bloom, petal fall stages and up to 5 weeks after petal fall. May be used up to 12 days of harvest. Do not feed treated hulls to dairy animals or animals being finished for slaughter.

APPLES Apply 30-60 lbs per acre to control Scab, Frogeye leaf spot, Bitter rot, Black rot, Botrysophaeria. No time limitation.

CHERRIES. Apply 40-100 lbs. per acre to control Brown rot, Cherry leaf spot. No time limitation.

CITRUS (Grapefruit, Lemons, Limes, Oranges) Apply 40-100 lbs. per acre to control Brown rot (Phytophthora). Do not feed citrus by-products to dairy animals or animals being finished for slaughter. No time limitation.

GRAPES Apply 20-50 lbs per acre to control Black rot, no time limitation.

NECTARINES, PEACHES: Apply 40-50 lbs per acre to control Brown rot, Scab, Coryneum blight. No time limitation.

PEARS. Apply 40-60 lbs. per acre to control Pear scab. Do not use on D'Anjou pear. No time limitation.

PLUMS, PRUNES Apply 40-50 lbs per acre to control Brown rot. No time limitation.

STRAWBERRIES Apply 30-38 lbs per acre to control Botrytis rot. Make first application when new growth starts. Repeat at weekly intervals up to harvest. No time limitation.

FIELD AND VEGETABLE CROPS

BEANS, PEAS Apply 40-60 lbs per acre pre-plant soil treatment to control Damp-off (Beans), Root rot. Work into top 3 or 4 inches of soil before planting.

CELERY Apply 20 lbs per acre for plant bed treatment to control Damp-off, Pink rot. Apply when all seeds have germinated and repeat when necessary.

COTTON Soil Treatment Apply 50-60 lbs per acre to control Soreshin. Apply in the seed furrow at planting time. 36" rows, apply 4 lbs per 1,000 feet. Coverage of material with soil is important.

CUCUMBERS, CANTALOUPES, SQUASH, PUMPKINS, HONEYDEW MELONS, WATERMELONS Apply 20-30 lbs per acre to control angular leaf spot, Anthracnose, Downy mildew. Use lower rates on young plants. Start applications when runners start to form or at first sign of disease. No time limitation.

POTATOES Apply 40-50 lbs per acre to control early and late blight. Use lower rates on young plants. Make first application when plants are 2-4 inches high or when disease is present. Repeat at 5-8 day intervals as needed. No time limitation.

POTATO SEED PIECE TREATMENT: Apply 1-1½ lbs per 100 lbs of seed piece to control Damp-off. Seed rot. Treat pieces within 6 hours after cutting. Store for 2 days in open crates if planting is delayed.

TOMATOES Apply 20-30 lbs per acre on young plants, 30-50 lbs on mature plants to control Anthracnose, Septoria leaf spot, Early and Late blight. No time limitation.

RQ CAPTAN MIXTURE
HA-9099

QUOM-E

ACCEPTED
DEC 4 1984
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended for the pesticide registered under EPA Reg No. 1202-107

NET CONTENTS _____ LBS

DISTRIBUTED BY PureGro
WEST SACRAMENTO, CALIFORNIA

LOT NO _____ 5/84 D.F.

BEST AVAILABLE COPY