

DIRECTIONS FOR USE

Apply when air is calm with an efficient dust applicator, securing complete coverage with a light film of dust. Begin at first sign of infection and repeat at 5-10 day intervals as needed to maintain control.

APPLES - Apply 30-80 lbs. per acre to control Scab, Frog-eye leaf spot, Bitter rot, Black rot, Botrytisphaeria. No time limitation.

APRICOTS - Apply 30-50 lbs. per acre to control Brown rot, (Twig blight), Jacket rot. No time limitation.

CHERRIES - Apply 40-100 lbs. per acre to control Brown rot, Cherry leaf spot. No time limitation.

CITRUS (Grapefruit, Lemons, Limes, Oranges) - Apply 40-100 lbs. per acre to control Brown rot (Phytophthora). Do not feed citrus by-products to dairy animals or animals being finished for slaughter. No time limitation.

GRAPES - Apply 20-50 lbs. per acre to control Black rot. No time limitation.

NECTARINES, PEACHES - Apply 40-50 lbs. per acre to control Brown rot, Scab, Coryneum blight. No time limitation.

PEARS - Apply 40-80 lbs. per acre to control Pear scab. Do not use on D'Anjou pear. No time limitation.

PLUMS, PRUNES - Apply 40-50 lbs. per acre to control Brown rot. No time limitation.

STRAWBERRIES - Apply 30-38 lbs. per acre to control Botrytis rot. Make first application when new growth starts. Repeat at weekly intervals up to harvest. No time limitation.

BEANS, PEAS - Apply 40-60 lbs. per acre pre-plant soil treatment to control Damp-off (Beans), Root rot. Work into top 3 or 4 inches of soil before planting.

CELERY - Apply 30-60 lbs. per acre for plant bed treatment to control Damp-off, Pink rot. Apply when all seeds have germinated and repeat when necessary.

(Continued on right panel)

Lot No. 24867RJW



**CAPTAN SULFUR
10-50 DUST**

ACTIVE INGREDIENT:

*Captan (N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide)	10.0%
Sulfur	50.0%
INERT INGREDIENTS:	40.0%
	100.0%

*U.S. Patent Numbers 2, 553, 770 and 2, 553, 771

**KEEP OUT OF RE/
OF CHILDREN**

CAUTION

Do not apply to cucurbits unless the variety is known to be sulfur tolerant under local conditions. Avoid drift to sulfur sensitive plants.

Avoid inhalation of dust. Avoid prolonged or repeated contact with skin. Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of any body of water.

Do not apply when weather conditions favor drift from the area.



BURN EMPTY CONTAINER
KEEP OUT OF SMOKE

Net Contents 50 Lbs.

(Continued from left panel)

COTTON - Soil treatment. Apply 50-60 lbs. per acre to control Soreshin. Apply in the seed furrow at planting time. 38" rows, apply 4 lbs. per 1,000 feet. Coverage of material with soil is important.

CUCUMBERS, CANTALOUPEs, SQUASH, PUMPKINS, HONEYDEW MELONS, WATERMELONS - Apply 20-30 lbs. per acre to control angular leaf spot, Anthracnose, Downy mildew. Use lower rates on young plants. Start applications when runners start to form or at first sign of disease. No time limitation.

POTATOES - Apply 40-50 lbs. per acre to control early and late blight. Use lower rates on young plants. Make first application when plants are 2-4 inches high or when disease is present. Repeat at 5-8 day intervals as needed. No time limitation.

POTATO SEED PIECE TREATMENT - Apply 1-1 1/2 lbs. per 100 lbs. of seed piece to control Damp-off, Seed rot. Treat pieces within 6 hours after cutting. Store for 2 days in open crates if planting is delayed.

TOMATOES - Apply 20-30 lbs. per acre on young plants, 30-50 lbs. on mature plants to control Anthracnose, Septoria leaf spot, Early and Late blight. No time limitation.

CONDITIONS OF SALE: 1. PureGro Company

CONDITIONS OF SALE: 1. PureGro Company (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond the manufacturer's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above) resulting from handling, storage, and use of this material.

USDA Reg. No. 1202-120

