



CHLORDANE 8 LIQUID

ACTIVE INGREDIENTS:

Technical Chlordane*73%
Petroleum Hydrocarbons21%

INERT INGREDIENTS:6%

*Equivalent to 43% Octachloro—4, 7—Methano-
tetrahydroindane and 30% Related Compounds.

Contains 8 lbs. of Technical Chlordane Per Gallon

WARNING

EPA Reg. No. 1202-123-AA

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs.



LOS ANGELES, CALIFORNIA


NET CONTENTS GALLON

SEE ADDITIONAL CAUTIONS 5TH PANEL

ACCEPTED

8/31/72

1202-123



CHLORDANE 8 LIQUID

Fruit Crops: Rate given in pints per 100 gallons water for ground sprayer unless stated. Aircraft use 5-30 gallons water per acre unless equivalent rate of Chlordane 8 Liquid.

APRICOTS, NECTARINES, PEACHES: Apply $\frac{1}{2}$ - $\frac{3}{4}$ pint per 100 gallons water to control catfacing insects, plum curculio. Several repeat applications may be necessary. Do not apply within 30 days of harvest. Do not exceed 6 pints per acre.

BLACKBERRIES, RASPBERRIES: Apply 2 pints per acre in 100-300 gallons water to control strawberry root weevil. Apply before fruit begins to form or after harvest.

CITRUS (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangerines, Tangelos): Apply 2 pints per 100 gallons water to control ants, grasshoppers, termites. Do not apply within 14 days of harvest. Apply to trunk and larger branches and to outer foliage only at drip line. For soil treatment apply when no fruit is present. Do not use over 10 pints per acre.

GRAPES: Apply 2-4 pints per acre in 100-400 gallons of water to control thrips. Do not apply after edible parts start to form.

Apply 3-10 pints per acre in 100-400 gallons water as a soil application to control Japanese beetle grubs. Apply when no fruit is present.

STRAWBERRIES: Apply 1-2 pints per acre in 100-200 gallons water to control crown borer, field crickets, lesser cornstalk borer, spittle bug, strawberry root worm (adult). Do not apply after fruit begins to form. Apply 10 pints per acre in 20-200 gallons water to control strawberry root weevil, white grubs, strawberry root worm, wireworm. Apply at a time of planting or when no fruit is present.

Vegetable, Field, Forage and Seed Crops: Rates given in pints per acre. To be applied by ground sprayer at 20-100 gallons water per acre unless specified. By aircraft, 5-30 gallons water per acre.

ASPARAGUS: Apply $\frac{1}{2}$ pint per acre to control asparagus beetles. Post cutting use only.

BEANS, BLACKEYED PEAS: Apply 1-2 pints per acre to control flea beetles, green stink bug. Beans—do not apply to green or snap beans after pods begin to form. Blackeyed peas—do not apply after pods start to form.

BEETS: Apply 1-2 pints per acre to control cutworms, flea beetle, armyworm. Do not apply after seedling stage if tops are to be used as food.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, COLLARDS, KALE, KOHLRABI: Apply 1-2 pints per acre to control cabbage worms, flea beetle, green stink bugs, harlequin bug. Do not apply after head or edible parts start to form. To control cabbage maggots, spray 1-2 pints in 100 gallons water per acre on newly set plants spraying around base of plant as soon as set in the field. Stir or wash the treatment into the soil.

CARROTS, CELERY: Apply 1-2 pints per acre to control carrot rust fly. Do not apply after plants are half grown or start to bunch on celery. No time limitation on carrots.

COTTON: Apply 1-3 $\frac{1}{2}$ pints per acre to control bollweevil, cotton fleahoppers, darkling beetle, fall armyworms, grasshoppers, tarnished plant bug, thrips, sand wireworms. Do not graze treated fields by dairy animals or animals being finished for slaughter.

Do not apply after bolls open.

EGGPLANT: Apply 1-2 pints per acre to control flea beetles, serpentine leaf miner. Do not apply after edible parts start to form.

LETTUCE: Apply $\frac{1}{2}$ -2 pints per acre to control armyworms, lygus bugs, serpentine leafminer, spotted cucumber beetle. Do not apply after edible parts start to form.

8/31/72

1202-123

ONIONS: Apply 1-2 pints per acre to control thrips. To control onion maggots, spray 2 pints per acre in a 6 inch band along onion row when onions are 1 inch out of the ground. Repeat in 7-10 days. Do not apply to spring or green onions.

PEAS: Apply 1-1½ pint per acre to control pea leafminers. Do not apply within 15 days of harvest. Do not feed treated forage to livestock.

PEPPERS: Apply 1½ - 2 pints per acre to control cutworms, mole crickets. Work into soil around base of plant. Do not make application after edible parts start to form. To control wireworms. Apply 3-10 pints per acre. Work into soil prior to or at planting.

POTATOES: Apply ¾ - 1½ pint per acre to control Colorado potato beetle, flea beetle, serpentine leafminers. No time limitation.

RICE: Apply 1½ - 3 pints per acre to control rice leafminers, armyworms. Do not apply after heads start to form. Do not graze treated forage. No restriction on use of grain.

SMALL GRAINS (Barley, Oats, Rye, Wheat): Apply 1-3 pints per acre to control armyworms. Do not apply after heads start to form. Do not graze treated forage. No restriction on use of grain.

TOMATOES: Apply 1-5 pints per acre to control flea beetles, serpentine leafminers, tomato hornworm. Do not apply after fruit begins to form

PREPLANTING SOIL APPLICATION OR AT TIME OF PLANTING OR TRANSPLANTING (Beans, Beets, Blackeyed Peas, Broccoli, Brussels Sprouts, Cabbage, Cantaloupes, Cauliflower, Celery, Collards, Corn, Cowpeas, Cucumbers, Eggplants, Garlic, Kale, Kohlrabi, Leeks, Lettuce, Melons, Okra, Onions, Peas, Peppers, Pimentos, Radishes, Rutabagas, Shallots, Squash (Summer), Squash (Winter), Sweet Potatoes, Tomatoes, Turnips, and Watermelons): Apply 9-10 pints per acre in 20-200 gallons water to control wireworms, white grubs, cutworms, seed corn maggot, diatritica larvae

WARNING

For protection of honey bees avoid use, if possible, during periods when bees are visiting the crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance.

This product is toxic to fish, birds, and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning (Keep out of smoke).

CONDITIONS OF SALE. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

LOT NO.

6 72 CN