

PURE GRO[®] SE

Use white mites and two-spotted mites dusting equipment. Do not apply to the soil. Apply to plant surfaces, treatable to tender and young fruit surfaces. Apply when mites first appear and repeat at 7-10 day intervals as needed to maintain control, but do not apply closer than 14 days.

FRUIT and VINE CROPS

APPLES, PINK: Brown two-spotted, Schoene, McDaniel, yellow (carmine), Willamette, Pacific and apple rust mites; for clean up of European red mite and clover mite -- Apply 40-50 lbs. per acre at petal fall to second cover or whenever mite population appears. Do not apply within 7 days of harvest.

CHERRY, NECTARINES, PEACHES, PLUMS, PRUNES: European red, clover (brown almond), two-spotted, Schoene, McDaniel, Pacific, Willamette, peach silver, plum nursery, and yellow mites -- Apply 40-50 lbs. per acre at petal fall to first cover or whenever mite population appears. If repeat application is necessary, wait 30 days to avoid excessive residue. Do not apply within 7 days of harvest on cherries, plums, and prunes; 14 days on peach, nectarine.

CITRUS (ORANGES, GRAPEFRUIT, LEMONS, LIMES, TANGERINES, TANGELOES, TANGERINES): Apply 50-100 lbs. per acre to control mites (Citrus red, Yuma, Citrus flat, six-spotted). Do not apply within 7 days of harvest.

PEACHES: Brown almond (clover), European, McDaniel, Pacific, two-spotted, Willamette and yellow mites -- Apply 10-35 lbs. per acre. Do not apply within 7 days of harvest.

RASPBERRIES, STRAWBERRIES, BLACKBERRIES, BOYSENBERRIES: Brown almond (clover), two-spotted, European red, Pacific, McDaniel, Willamette and yellow mites -- Apply 15-30 lbs. per acre. Do not apply within 2 days of harvest.

STRAWBERRIES: Cyclamen, two-spotted, strawberry (Atlantic) and six-spotted mites -- Apply 25-35 lbs. per acre. Do not apply within 7 days of harvest.

WALNUTS, FILBERTS: Brown almond (clover), two-spotted, European red, Pacific, McDaniel, Willamette and yellow mites -- Apply 40 lbs. per acre. Do not apply within 14 days of harvest. Do not feed hulls to livestock.

Vegetable and Field Crops

ALFALFA, CLOVER (Seed Crops): Two-spotted, Strawberry (Atlantic), Carmine, and Pacific mites -- Apply 25-35 lbs. per acre. Use restricted to California and to Grant and Adams Counties, Washington. Fields not to be grazed by livestock. Crop cuttings or residues not to be fed or baled. The fields must be cleared after harvest by burning or discing into the soil.

CUCUMBERS, CANTALOUPES, SQUASH: Apply 20-35 lbs. per acre to control two-spotted and Pacific mites. Do not apply within 7 days of harvest.



KELTHANE[®] 4 DUST

AN AGRICULTURAL MITICIDE

ACTIVE INGREDIENT
1,1-bis (p-chlorophenyl)-2,2,2-Tri-Elongethane . . . 4.0
INERT INGREDIENTS 96.0

lbs./

*Registered trademark of Rohm & Haas Company

E.P.A. Reg. No. 1.0-116-AA

CAUTION

KEEP OUT OF REACH OF CHILDREN

Keep out of reach of children. Avoid inhalation, contact with eyes, skin, and clothing. Wash thoroughly after handling.

WEAR THE FOLLOWING EQUIPMENT:

Observe the limitations given on the label for species treated. Do not use on feed or forage to be fed to dairy animals or animals being finished for slaughter.

This product is toxic to fish. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

The Federal Food and Drug Administration has set off maximum tolerances for certain pesticide residues. In order that residues on these agricultural crops will not exceed established tolerances, only at recommended rates, and do not apply closer to harvest than specified.

1. WILLOW, STRAWBERRY (ATLANTIC): 20-35 lbs. per acre. Do not apply within 7 days of harvest.

2. WILLOW, STRAWBERRY (ATLANTIC): 15-35 lbs. per acre depending on days of harvest. Do not graze.

3. SPOTS: Two-spotted mites -- Apply 20-35 lbs. for mid to late season.

4. LEAVES, EGGPLANT, PEPPERS: Pacific spotted mites -- Apply 20-35 lbs. per acre. Do not apply within 7 days of harvest.

5. ORNAMENTALS, FLOWERS, NURSERY STOCK: Greenhouse or outdoor plants, make present and repeat as necessary.

EXCLUSIONS OF SALE: 1. PureGro material conforms to the chemical definition for use as directed herein. Manufacturer or representative to make, and merchantability, guarantee or represent this material.

2. Critical and unforeseeable factors from eliminating all risks in commercial use include, but are not limited to the material is applied, lack of control on other plants or crops. Such risks reasonably fit for the uses stated followed. Buyer and user acknowledge those assumed by the manufacturer upon use of this material.



10-116-AA

11-19-72
ACCEPTED WITH COMMENTS



KELTHANE® 4 DUST

AN AERIAL SPRAY MITICIDE

ACTIVE INGREDIENT	
1,1-Ets- <i>p</i> -chlorophenyl)-2,2,2-trichloroethanol ... 4.7	
INERT INGREDIENTS	95.3
	100.0

*Registered trademark of Rohm & Haas Company

E.P.A. Reg. No. 120z-126-AA

CAUTION

KEEP OUT OF REACH OF CHILDREN

Adults: Eat fruit. Avoid inhalation of dust. Avoid ingestion of dust and wash off. Wash face with soap and water.

REHEAT THE FOLLOWING INFORMATION

Follow all limitations given on the label for specific rates, do not use on feed or forage to be fed to dairy animals or animal to be finished for slaughter.

This product is toxic to fish. Keep out of any body of water. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment, tools, pose, or wastes.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues in feed and forage crops will not exceed established tolerances, only at recommended rates, and do not apply closer to harvest than specified.

Do not apply to:

LAWNS: Twospotted, strawberry, Atlantic, Pacific, two-spotted mites -- Apply 15-30 lbs. per acre. Do not graze treated vines until after harvest. Do not apply within 7 days of harvest.

FRUIT: Atlantic (strawberry), Pacific, two-spotted and desert mites -- Apply 15-35 lbs. per acre depending on size of plants. Do not apply within 14 days of harvest. Do not graze treated fields by eat or dairy cattle.

CROPS: Two-spotted mites -- Apply 15-30 lbs. per acre during early season. 20-35 lbs. for mid to late season. Do not apply within 7 days of harvest.

VEGETABLES, TOMATOES, EGGPLANT, PEPPERS: Pacific, tropical, tomato russet and two-spotted mites -- Apply 20-35 lbs. per acre. Do not apply within 14 days of harvest.

PORVENEMENTS, FLOWERS, NURSERY STOCK AND SHADE TREES: For control of mites in greenhouse or outdoor plants, make full coverage applications when stated above present and repeat as necessary.

NOTICES OF SALE: 1. PureGro Company (manufacturer) warrants that the material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS, or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond the manufacturer's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated herein and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above) resulting from handling, use, and use of this material.

TURN EMPTY CONTAINER
KEEP OUT OF SUN

Lot No.

12-12-72



May 11, 1972