

DIRECTIONS FOR USE

Field Crops and Vegetables

ASPARAGUS: Apply 20-40 lbs. per acre to control asparagus beetle. Apply during cutting season. Do not repeat application within 3 days. Do not apply 1 day before harvest.

BEANS: Mexican bean beetle, leafhoppers, bean leaf beetle, cucumber beetles, Western bean cutworm, flea beetles, Japanese beetle, velvet-bean caterpillar, armyworms, tarnished plant bug, corn earworm. Apply 20-40 lbs. per acre. No time limitation.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI: Flea beetle, harlequin bug, corn earworm, armyworms. For small loopers after heads form, use SEVIN every 5 to 7 days. Apply 20-40 lbs. per acre. Do not apply 3 days before harvest.

CORN: Corn earworm, (Apply to silks as soon as they appear. Make 3 to 4 applications at 2-3 day intervals. Re-apply immediately after rain.), corn rootworm adults, European corn borer, fall armyworm, sap beetles, Japanese beetle, flea beetles, leafhoppers -- Apply 20-40 lbs. per acre. No time limitation.

COTTON: Thrips, flea beetle, fleahopper, leafworm, striped blister beetle, boll weevil, bollworm, fall armyworm, leaf perforator, leaf rollers, leafhoppers, tarnished plant bug, light to moderate numbers of Western lygus bugs, (aphids will be repressed by repeated applications in a schedule), pink bollworm, salt marsh caterpillar, stink bugs, grasshoppers -- Apply 30-40 lbs. per acre. No time limitations. Allow 7 days before grazing or harvesting crop residue for feed or bedding for dairy or meat animals.

LEAFY VEGETABLES AND ROOT CROPS: Chinese cabbage, collards, dandelion, Endive, lettuce, leaf lettuce, mustard greens, Parsley, Spinach, Swiss chard) -- Apply 1-10 lbs. per acre to control flea beetles, leafhoppers, harlequin bug, six-spotted leafhopper (aster yellows vector), corn earworm, armyworms, lygus and stink bugs. Do not apply within 14 days of harvest.

PEAS: Leafhoppers, colored potato beetle, armyworm, grasshoppers -- Apply 1-10 lbs. per acre. No time limitation.

POTATO: Colorado potato beetle, flea beetles, leafhoppers -- Apply 20-40 lbs. per acre. Foliage application only.

SUGAR BEETS: Leafhoppers, armyworms, webworms, grasshoppers -- Apply 20-40 lbs. per acre. Do not apply 10 days before harvest.

TOMATO, EGGPLANT, PEPPER: Tomato fruitworm, fall armyworm, tomato hornworm, European corn borer, lace bugs, Colorado potato beetle, flea beetles, leafhoppers -- Apply 20-80 lbs. per acre. No time limitation.

CUCUMBERS, SQUASH, PUMPKIN, MELONS: Flickerworm, melonworm, cucumber beetles, squash bug, flea beetles, leafhoppers -- Apply 20 lbs. per acre. No time limitation.

HEAD LETTUCE, HORSERADISH, PARSNIPS, RADISHES, RUTABAGAS: Apply 20-40 lbs. per acre to control flea beetles, leafhoppers, harlequin bugs, six-spotted leafhopper (aster yellows vector), corn earworm, armyworms, lygus bug, stink bug. Do not apply within 3 days of harvest.

BEETS, SWISS CHARD, TURNIPS: Apply 20-40 lbs. per acre to control flea beetles, leafhoppers, harlequin bugs, six-spotted leafhopper (aster yellows vector), corn earworm, armyworms, lygus bug, stink bug. Do not apply within 3 days of harvest. Do not apply within 14 days of harvest if tops are to be used as food or feed.



## SEVIN® SULFUR DUST 5-50

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methyl-carbamate) . . . . .	5.0%
Sulfur . . . . .	50.0%
INERT INGREDIENTS . . . . .	45.0%
	100.0%

Fruit and Nut Crops

For codling moth, orange tortrix, and every 10 to 14 days.

APPLES, PEARS: Codling moth, California pear slug, harvest.

CHERRY, PLUM, PRUNE: Orange tortrix, Webworm, apply 1 day before harvest.

GRAPES: Grape leaf moth, Japanese beetle per acre. No time limitation.

PEACHES, NECTARINES: Orange tortrix, tussock, moth per acre. Do not apply for nectarines.

Forage and Feed Crops

ALFALFA, CLOVERS, BEANS: Thrips, leafhoppers, beetle, Japanese beetle, cloverworm, armyworm per acre. No time limitation.

FORAGE GRASSES, PASTURES: For optimum thrip control, apply up to half a pound grain and insect dust into feed (removed from boot.) 3 days before harvest.

WHEAT, BARLEY, GRAIN: Direct dust into feed (removed from boot.) 3 days before harvest.

CONDITION OF SALE: Application, weather not specifically required for this product are beyond the control of use, storage or directions given here.

E.P.A. REG. NO. 1202-139-AA

Registered trademark of Union Carbide Corporation for the active ingredient (Carbaryl).

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin; wash thoroughly after using. Avoid storage near feed and food products.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

DO NOT USE  
ON EMPTY CONTAINER  
KEEP OUT OF SMOKE



PESTICIDE ACT  
ON REGISTER  
2-139



## SEVIN® SULFUR DUST 5-50

### ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methyl-carbamate) . . . . .	5.0%
Sulfur . . . . .	50.0%
INERT INGREDIENTS . . . . .	45.0% 100.0%

E.P.A. REG. NO. 1202-139-AA

Registered trademark of Union Carbide Corporation  
for the active ingredient (Carbaryl).

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin; wash thoroughly after using. Avoid storage near feed and food products.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

### Fruit and Nut Crops

For codling moth, oriental fruit moth and plum curculio. Apply at petal fall and every 10 to 14 days thereafter until control is achieved.

APPLES, PEARS: Codling moth, green apple aphid, pear psylla, orange tortrix, California pear slug. Apply 100-200 lbs. per acre. Do not apply 1 day before harvest.

CHERRY, PLUM, PRUNE: Cherry fruit fly, fruit tree leaf roller, peach twig borer, orange tortrix, Western tussock moth -- Apply 80-120 lbs. per acre. Do not apply 1 day before harvest.

GRAPES: Grape leaf folder, grape leafhopper, grape leaf skeletonizer, grape berry moth, Japanese beetle, June beetles, red-banded leaf roller -- Apply 40-60 lbs. per acre. No time limitation.

PEACHES, NECTARINES: Oriental fruit moth, peach twig borer, codling moth, orange tortrix, tussock moth, fruit tree leaf roller, Japanese beetle. Apply 80-160 lbs. per acre. Do not apply 1 day before harvest for peaches and 3 days before harvest for nectarines.

### Forage and Feed Crops

ALFALFA, CLOVERS, PEANUTS, SOYBEANS, COWPEAS: Blister beetles, Mexican bean beetle, thrips, leafhoppers, three-cornered alfalfa hopper, cucumber beetles, bean leaf beetle, Japanese beetle, alfalfa caterpillar, velvetbean caterpillar and green cloverworm, armyworms, webworms, corn earworm, grasshoppers. Apply 20-30 lbs. per acre. No time limitation.

FORAGE GRASSES, PASTURE: Apply 25-30 lbs. per acre to control armyworms, thrips. For optimum thrip control in grasses grown for seed, direct dust into boot. May be applied up to harvest.

SORGHUM (Grain and Forage): Armyworm, webworm, corn earworm, sorghum midge. Direct dust into forming heads. Apply three or four days after heads have emerged from boot., grasshoppers -- Apply 30-40 lbs. per acre. Do not apply 1 day before harvest of grain. No time limitation on sorghum forage.

**CONDITIONS OF SALE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

BURN EMPTY CONTAINER  
KEEP OUT OF SMOKE



LOS ANGELES, CALIFORNIA

NET WT 15 LBS.

LBS.