

STRAWBERRIES: Apply 1/2 pint per acre to control aphid, mites. Do not apply within 14 days of harvest.

SUGAR BEETS: Apply 1/4 pint per acre to control aphids, mites, army worms, Sugar beet webworm, grasshoppers, Climbing worms. Do not apply within 20 days of harvest. If tops are to be fed to livestock, do not apply within 60 days of harvest.

TOMATOES: Apply 1/2 pint per acre to control aphids, army worms, flea beetles, grasshoppers, mites. Do not apply within 15 days of harvest.

TURNIPS: Control areas of cutworm, cabbage aphid. Apply 1/2 pint per acre. Do not apply within 15 days of harvest. 20 days if tops are to be used as food or feed.

VETCH: Apply 1/2 pint per acre to control spotted alfalfa aphid. Do not apply within 15 days of harvest.

SMALL GRAINS (Barley, Oats, Wheat): Apply 1/2 pint per acre to control aphid, black grass bug, brown wheat mite, pink bug. Do not apply within 15 days of harvest.

WARNING!


Dangerous if Swallowed, Inhaled, or Absorbed Through Skin. Rapidly Absorbed Through Skin! Do not get in eyes, on skin, or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for parathion protection. Keep away from protected persons, out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms, and face thoroughly with soap and water before eating or drinking. Wash all contaminated clothing with soap and water before reuse.

Do not use this toxin to fish and wildlife. If as a gutter wash, treated areas may be harmed. Keep out of lakes, ponds, streams. Do not apply to any area not specified on the label.

Use equipment and containers and dispose of wastes by burying in deep land away from water supplies. Containers should be disposed of by puncturing holes in top and bottom and washing with water. Do not reuse. Keep out of reach of children.

E-10-N

P10966R1W



DIRECTIONS FOR USE

SEVIN® DUST 7.5

Field Crops and Vegetables

COTTON: Thrips, flea beetle, fleahopper, leafworm, striped blister beetle, boll weevil, bollworm, fall armyworm, leaf perforator, leaf rollers, leafhoppers, tarnished plant bug, light to moderate numbers of V. stem Lygus bugs, (aphids will be repressed by repeat applications in a schedule), pink bollworm, salt marsh caterpillar, stink bugs, grasshoppers. Apply 20 lbs. per acre. No time limitations. Allow 7 days before grazing or harvesting crop residue for feed or bedding for dairy or meat animals.

BEANS: Mexican bean beetle, leafhoppers, bean leaf beetle, cucumber beetles, Western bean cutworm, flea beetles, Japanese beetle, velvet-bean caterpillar, corn earworm, armyworms, stink bugs, tarnished plant bug (lima bean pod borer) corn earworm, Lygus and stink bugs--Apply 15-25 lbs. per acre. No time limitation.

PEAS: Leafhoppers, Colorado potato beetle, armyworms, grasshoppers--Apply 15-20 lbs. per acre. No time limitation.

CORN: Corn earworm, (Apply to silks as soon as they appear. Make 3 to 4 application at 2-3 day intervals. Re-apply immediately after rain.), corn rootworm adults, European corn borer, fall armyworm, sap beetles, Japanese beetle, flea beetles, leafhoppers--Apply 15-25 lbs. per acre. No time limitation.

SUGAR BEETS: Leafhoppers, armyworms, webworms, grasshoppers--Apply 15-25 lbs. per acre. Do not apply 14 days before harvest

POTATO: Colorado potato beetle, flea beetles, leafhoppers--Apply 15-25 lbs. per acre. Foliage application only.

TOMATO, EGGPLANT, PEPPER: Tomato fruitworm, fall armyworm, tomato hornworm, European corn borer, stink bugs, lace bugs, Lygus bugs, Colorado potato beetle, flea beetles, leafhoppers--Apply 15-50 lbs. per acre. No time limitation.

CUCUMBERS, SQUASH, PUMPKIN, MELONS: Pickleworm, melonworm, cucumber beetles, squash bug, flea beetles, leafhoppers--Apply 15-25 lbs. per acre. No time limitation.

ASPARAGUS: Apply 15-25 lbs. per acre to control asparagus beetle. Apply during cutting season. Do not repeat application within 3 days. Do not apply 1 day before harvest.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI: Flea beetle, harlequin bug, corn earworm, armyworms. For small loopers after heads form, use SEVIN every 5 to 7 days--Apply 15-25 lbs. per acre. Do not apply 3 days before harvest.

ACTIVE INGREDIENTS:
Carbaryl (1-naphthyl N-methylcarbamate) 7.5%
INERT INGREDIENTS 92.5%
KEEP OUT OF REACH OF CHILDREN. 100.0%

CAUTION

Harmful if swallowed. Avoid breathing dust. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid storage near feed and food products.

This product is toxic to fish. Keep out of lakes, ponds and streams.

Do not apply to any area not specified on the label.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.



Registered trademark of Union Carbide Corporation for the active ingredient (Carbaryl).

BURN EMPTY CONTAINER
KEEP OUT OF SMOKE

Fruit and Nut Crops

APPLES, PEARS: Codling moth, green apple aphid, pear psylla, orange tortrix, California pear slug. Apply 65-130 lbs. per acre. Do not apply 1 day before harvest.

PEACHES, APRICOTS, NECTARINES: Oriental fruit moth, peach twig borer, codling moth, orange tortrix, tussock moth, fruit tree leaf roller, periodical cicada, Japanese beetle. Apply 50-130 lbs. per acre. Do not apply 1 day before harvest for peaches and 3 days before harvest for apricots and nectarines.

GRAPES: Grape leaf folder, grape leafhopper, grape leaf skeletonizer, grape berry moth, Japanese beetle, June beetles, red-banded leaf roller--Apply 20-40 lbs. per acre. No time limitation.

CHERRY, PLUM, PRUNE: Cherry fruit fly, fruit tree leaf roller, peach twig borer, orange tortrix, Western tussock moth--Apply 50-80 lbs. per acre. Do not apply 1 day before harvest.

FILBERT: Filbert worm, filbert aphid, filbert leaf roller--Apply 65 lbs. per acre. No time limitation.

Forage and Feed Crops

ALFALFA, CLOVERS, PEANUTS, SOYBEANS, COWPEAS: Blister beetles, Mexican bean beetle, thrips, leafhoppers, three-cornered alfalfa hopper, cucumber beetles, bean leaf beetle, Japanese beetle, alfalfa caterpillar, velvetbean caterpillar and green cloverworm armyworms, webworms, corn earworm, stink bugs, grasshoppers. Apply 15-20 lbs. per acre. No time limitation.

SORGHUM (Grain and Forage): Armyworm, webworms, corn earworm, stink bugs, sorghum midge, grasshoppers--Apply 20-25 lbs. per acre. Do not apply 21 days before harvest of grain. No time limitation on sorghum forage.


RICE, FORAGE GRASSES, PASTURE: Armyworms, stink bugs, thrips. Apply 15 to 20 lbs. per acre. For optimum thrip control in grasses grown for seed, direct dust into boot. Do not apply to Rice within 14 days of harvest. May be applied up to harvest on Pastures and Forage Grasses.

NONWARRANTY

Because the use of this material is subject to conditions beyond their control, neither the Pacific Guano Company, nor any other Seller hereof warrants, either expressed or implied, the use thereof or the results of such use, whether or not used in accordance with directions. Neither of the foregoing will or shall be responsible for the results of any use of this material.

Lot No. _____

NET CONTENTS _____



USDA 16-750 _____