

DIRECTIONS FOR USE

APPLES, PEARS: Apply 30-100 lbs. per acre to control green apple aphid, woolly apple aphid, San Jose scale crawlers, mites, pear psylla. Slight russeting or other phytotoxic effects may occur on some varieties of apples such as Golden Delicious. Do not apply within 14 days of harvest.

CHERRIES: Apply 30-100 lbs. per acre to control black cherry aphids, cherry fruit flies, San Jose scale crawlers. Do not apply within 10 days of harvest.

EGGS: Apply 30-80 lbs. per acre to control vinegar flies, dried fruit beetles. Do not apply within 5 days of harvest.

GRAPES: Apply 30 lbs. per acre to control Pacific spider mites, leafhopper, grape berry moth, grape leaf folder. Do not apply within 10 days of harvest.

LEMONS, ORANGES: Apply 100-250 lbs. per acre to control citrus aphid, brown soft scale crawlers. Do not apply within 21 days of harvest.

NECTARINES, PEACHES, APRICOTS: Apply 40-130 lbs. per acre to control aphid, brown mites, two spotted mites, San Jose scale crawlers, olive scale crawlers. Do not apply within 20 days of harvest on peaches, 10 days on apricots and nectarines.

PLUMS, PRUNES: Apply 40-130 lbs. per acre to control leaf curl plum aphid, mealy plum aphid, thistle aphid, brown mites, European red mites. Do not apply within 10 days of harvest.

STRAWBERRIES: Apply 20-40 lbs. per acre to control cyclamen mites, aphids, two spotted mite. Do not apply within 5 days of harvest.

WALNUTS: Apply 40-150 lbs. per acre to control aphids, mites, scale crawlers. Do not apply after shucks split.

ALMONDS: Apply 40-150 lbs. per acre to control mites. Do not apply within 1 day of harvest.

BEANS (SNAP): Apply 20-40 lbs. per acre to control bean aphid, leaf miners, Mexican bean beetles, banded cucumber beetles. Do not apply within 7 days of harvest.

BEETS, LETTUCE, SPINACH: Apply 15 lbs. per acre to control aphids, dipterous leaf miners. Do not apply within 14 days of harvest.

BROCCOLI, CABBAGE, CAULIFLOWER, KALE: Apply 30-50 lbs. per acre to control diamondback moths, aphids, imported cabbageworm. Do not apply within 7 days of harvest except kale 10 days before harvest.

(Continued on right panel)

Lot No. _____ R27066R JW



**DIAZINON[®] SULFUR
3-25 DUST**

ACTIVE INGREDIENTS:	
O,O-diethyl O-(2-isopropyl-4-methyl 6-pyrimidinyl) phosphorothioate	3.0%
Sulfur	25.0%
INERT INGREDIENTS	72.0%
	100.0%

© Reg. F. M. Ceigy Chemical Co.

KEEP OUT OF REACH OF CHILDREN.


CAUTION

Harmful if swallowed. May be absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid storage near food and feed products.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label.

Do not apply when weather conditions favor drift of dust from areas treated.

Containers should be disposed by burning (Keep out of smoke).



NET WEIGHT 50 lbs.

(Continued from left panel)

CANTALOUPE, CUCUMBERS, MUSKMELONS, SQUASH, WATERMELONS, PEPPERS: Apply 40-45 lbs. per acre to control aphids, leafhoppers, mites, thrips, serpentine leaf miners. Do not apply within 7 days of harvest on squash, cucumbers, peppers; 3 days on cantaloupe, muskmelons, watermelons.

CELERY: Apply 15 lbs. per acre to control aphids, flea beetle. Do not apply within 10 days of harvest.

ONIONS: Apply 30-40 lbs. per acre to control thrips. Do not apply within 10 days of harvest.

POTATOES: Apply 30-65 lbs. per acre to control aphid, flea beetle, Colorado potato beetle, dipterous leaf miners, banded cucumber beetle, leafhoppers, southern armyworms. Do not apply within 11 days of harvest. Foliage application.

TOMATOES: Apply 20-50 lbs. per acre to control aphid, dipterous leaf miner, banded cucumber beetles, fall armyworms, southern armyworms, vinegar fly. Do not apply within 3 days of harvest.

ALFALFA, CLOVER: Apply 20 lbs. per acre to control spotted alfalfa aphid, yellow clover aphid, pea aphid, leafhoppers, plant bug nymphs, mites. Do not apply within 7 days of cutting or grazing by animals being finished for slaughter.

CORN: Apply 40-50 lbs. per acre to control grasshoppers, corn leaf aphid, corn earworms (Apply when silks first appear. Two or three applications at two or three day intervals will be necessary). Do not apply within 2 days of harvest. Do not feed treated forage to animals being finished for slaughter within 2 days after treatment.

COTTON: Apply 20-30 lbs. per acre to control spider mites, aphids, leafhoppers, cotton leaf worm, cotton leaf perforator, lygus, salt marsh caterpillar. Do not apply within 14 days of harvest. Do not graze dairy or meat animals. Do not feed gin waste to livestock.

NONWARRANTY

Because the use of this material is subject to conditions beyond their control, neither the Pacific Guano Company, nor any other Seller hereof warrants, either expressed or implied, the use thereof or the results of such use, whether or not used in accordance with directions. Neither of the foregoing will or shall be responsible for the results of any use of this material.

U.S.D.A. Reg. No. _____