

DIRECTIONS FOR USE

APPLES, PEARS: Apply 30-100 lbs. per acre to control green apple aphid, woolly apple aphid, San Jose scale crawlers, mites, pear psylla. Slight russeting or other phytotoxic effects may occur on some varieties of apples such as Golden Delicious. Do not apply within 14 days of harvest.

CHERRIES: Apply 30-100 lbs. per acre to control black cherry aphids, cherry fruit flies, San Jose scale crawlers. Do not apply within 10 days of harvest.

FIGS: Apply 25-60 lbs. per acre to control vinegar flies, dried fruit beetles. Do not apply within 5 days of harvest.

GRAPES: Apply 25 lbs. per acre to control Pacific spider mites, leafhopper, grape berry moth, grape leaf folder. Do not apply within 10 days of harvest.

LEMONS, ORANGES: Apply 70-200 lbs. per acre to control citrus aphid, brown soft scale crawlers. Do not apply within 21 days of harvest.

NECTARINES, PEACHES, APRICOTS: Apply 30-100 lbs. per acre to control aphid, brown mites, two spotted mites, San Jose scale crawlers, olive scale crawlers. Do not apply within 20 days of harvest on peaches, 10 days on apricots and nectarines.

PLUMS, PRUNES: Apply 30-100 lbs. per acre to control leaf curl plum aphid, mealy plum aphids, thistle aphid, brown mites, European red mites. Do not apply within 10 days of harvest.

STRAWBERRIES: Apply 20-30 lbs. per acre to control cyclamen mites, aphids, two spotted mite. Do not apply within 5 days of harvest.

WALNUTS: Apply 30-125 lbs. per acre to control aphids, mites, scale crawlers. Do not apply after shucks split.

ALMONDS: Apply 40-150 lbs. per acre to control mites. Do not apply within 1 day of harvest.

BEANS (SNAP): Apply 20-35 lbs. per acre to control bean aphid, leaf miners, Mexican bean beetles, banded cucumber beetles. Do not apply within 7 days of harvest.

BROCCOLI, CABBAGE, CAULIFLOWER, KALE: Apply 25-40 lbs. per acre to control diamondback moths, aphids, imported cabbageworm. Do not apply within 7 days of harvest except kale 10 days before harvest.

CARROTS: Apply 12 lbs. per acre to control aphids. Do not apply within 10 days of harvest.

ENDIVE, RADISHES: Apply 12 lbs. per acre to control aphids, dipterous leaf miners. Do not apply within 10 days of harvest.

(Continued on right panel)



DIAZINON® DUST 4

ACTIVE INGREDIENTS:

O,O-diethyl O-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate	4.0%
INERT INGREDIENTS:	96.0%
	100.0%

*Reg. T. M. Geigy Chemical Co.

CAUTION

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed. May be absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid storage near food and feed products.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label.

Do not apply when weather conditions favor drift of dust from areas treated.

Containers should be disposed by burning (Keep out of smoke).



U.S. D.A. Reg. No. 1202-152

Lot No. R13967R

ACCEPTED

NOV 13 1967

UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND PLANT GROWTH REGULATING ACT FOR ECONOMIC PEST CONTROL REGISTERED UNDER NO. 1202-152 SUBJECT TO ATTACHED COMMENTS

NET WEIGHT 50 lbs.

(Continued from left panel)

CANTALOUPE, CUCUMBERS, MUSKMELONS, SQUASH, WATERMELONS, PEPPERS: Apply 30-35 lbs. per acre to control aphids, leafhoppers, mites, thrips, serpentine leaf miners. Do not apply within 7 days of harvest on squash, cucumbers, peppers; 3 days on cantaloupes, muskmelons, watermelons.

ONIONS: Apply 20-30 lbs. per acre to control thrips. Do not apply within 10 days of harvest.

POTATOES: Apply 20-50 lbs. per acre to control aphids, flea beetles, Colorado potato beetle, dipterous leaf miners, banded cucumber beetles, leafhoppers, southern armyworms. Do not apply within 14 days of harvest. Foliage application.

TOMATOES: Apply 20-40 lbs. per acre to control aphids, dipterous leaf miners, banded cucumber beetles, fall armyworms, southern armyworms, vinegar fly. Do not apply within 3 days of harvest.

ALFALFA, CLOVER: Apply 15 lbs. per acre to control spotted alfalfa aphids, yellow clover aphids, pea aphids, leafhoppers, plant bug nymphs, mites. Do not apply within 7 days of cutting or grazing by animals being finished for slaughter.

CORN: Apply 30-40 lbs. per acre to control grasshoppers, corn leaf aphid, corn earworms. (Apply when silks first appear. Two or three applications at two or three days intervals will be necessary.) Do not apply within 2 days of harvest. Do not feed forage to livestock within 10 days following last application.

COTTON: Apply 20-25 lbs. per acre to control spider mites, aphids, leafhoppers, cotton leaf worms, cotton leaf perforators, lygus, salt marsh caterpillars. Do not apply within 14 days of harvest. Do not graze dairy or meat animals in treated fields. Do not feed gin waste to livestock.

CONDITIONS OF SALE: 1. PureGro Company (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond the manufacturer's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above) resulting from handling, storage, and use of this material.

ACCEPTED WITH COMMENTS