

1203-152
1/1/85
1/1/82

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Hazard: If swallowed. May be absorbed through skin. Do not breathe dust or spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling and before smoking or eating. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans or household pets. Do not permit children or pets to go onto sprayed areas until spray has completely dried.

Statement of Practical Treatment

If swallowed, call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled, remove victim to fresh air and apply artificial respiration if indicated.

If on skin, wash promptly with soap and water. Rinse thoroughly.

If in eyes, rinse with plenty of water and call a physician immediately.

Note to Physician: Atropine sulfate and 2-PPM should only be used when signs of cholinesterase inhibition are present.

LOS ANGELES POISON CONTROL CENTER (213) 664-2121 ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes, and estuaries.

Do not apply to water that will be used for recreational purposes and human and livestock consumption. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources.

Apply this product only as specified on this label. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity. In case of spill or accident, call CHEMTREC (800) 424-9300 day or night for assistance.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

REENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until dusts have settled.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written or oral warnings must include the following information:

CAUTION: Area to be treated with diazinon on (date of application). Do not enter without appropriate protective clothing until dusts have settled. If accidental exposure, consult the Statement of Practical Treatment portion of the pesticide label for first aid and medical attention recommendations.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store only in original container in locked storage area. If a spill occurs, refer to the pesticide disposal section for disposal. Avoid cross-contamination with other pesticides and fertilizers during handling and storage. Do not store near heat or open flame. Not for Use or Storage in or Around the Home.

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

PUREGRD DIAZINON DUST 4

ACTIVE INGREDIENTS:

0,0-diethyl 0-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate 33%

INERT INGREDIENTS:

96.0%
100.0%

EPA Reg. No. 1202-152-AA

Est. No.

ACCEPTED

SEP 17 1985

KEEP OUT OF REACH OF CHILDREN

Under the Federal Insecticide, Fungicide, and Rodenticide Act, this pesticide is registered under EPA Reg. No. 1202-152

CAUTION

SEE LEFT SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT.

CROP RECOMMENDATIONS

APPLES, PEARS: Apply 30-10 lbs. per acre to control green apple aphid, woolly apple aphid, San Jose scale crawlers, mites, pear psylla. Slight russetting or other phytotoxic effects may occur on some varieties of apples such as Golden Delicious. Do not apply within 14 days of harvest.

CHERRIES: Apply 30-100 lbs. per acre to control black cherry aphids, cherry fruit flies, San Jose scale crawlers. Do not apply within 10 days of harvest.

FIGS: Apply 25-60 lbs. per acre to control vinegar flies, dried fruit beetles. Do not apply within 5 days of harvest.

GRAPES: Apply 25 lbs. per acre to control Pacific spider mites, leafhopper, grape berry moth, grape leaf folder. Do not apply within 10 days of harvest.

LEMONS, ORANGES: Apply 70-200 lbs. per acre to control citrus aphid, brown soft scale crawlers. Do not apply within 21 days of harvest.

NECTARINES, PEACHES, APRICOTS: Apply 30-100 lbs. per acre to control aphid, brown mites, two spotted mites, San Jose scale crawlers, olive scale crawlers. Do not apply within 20 days of harvest on peaches, 10 days on apricots and nectarines.

Distributed by



WEST SACRAMENTO, CA 95691

NET CONTENTS _____ LBS. LOT NO. _____ 8/84 C.D.

IST 4

4.0%

96.0%

100.0%

CHILDREN

OXARY STATEMENT.

control green
cale crawlers,
toxic effects
Golden Delici-

of black cherry
s. Do not apply

gar flies, dried
rvest.
Pacific spider
' folder. Do not

to control
y within 21

ly 30-100 lbs.
ted mites, San
ot apply within
apricots and

DT NO.

MM G.D.

PLUMS, PRUNES: Apply 30-100 lbs. per acre to control leaf curl plum aphid, mealy plum aphids, thistle aphid, brown mites, European red mites. Do not apply within 10 days of harvest.

STRAWBERRIES: Apply 20-30 lbs. per acre to control cyclamen mites, aphids, two spotted mite. Do not apply within 5 days of harvest.

WALNUTS: Apply 30-125 lbs. per acre to control aphids, mites, scale crawlers. Do not apply after shucks split.

ALMONDS: Apply 40-150 lbs. per acre to control mites. Do not apply within 1 day of harvest.

BEANS (Snap): Apply 20-35 lbs. per acre to control bean aphid, leaf miners, Mexican bean beetles, banded cucumber beetles. Do not apply within 7 days of harvest.

BROCCOLI, CABBAGE, CAULIFLOWER, KALE: Apply 25-40 lbs. per acre to control diamondback moths, aphids, imported cabbageworm. Do not apply within 7 days of harvest. ~~Do not apply within 10 days of harvest on KALE.~~

CARROTS: Apply 12 lbs. per acre to control aphids. Do not apply within 10 days of harvest.

ENDIVE, RADISHES: Apply 12 lbs. per acre to control aphids, dipterous leaf miners. Do not apply within 10 days of harvest.

CANTALOUPE, CUCUMBERS, MUSKMELONS, SQUASH, WATERMELONS, PEPPERS: Apply 30-35 lbs. per acre to control aphids, leafhoppers, mites, thrips, serpentine leaf miners. Do not apply within 7 days of harvest on squash, cucumbers, peppers; 3 days on cantaloupes, muskmelons, watermelons.

ONIONS: Apply 20-30 lbs. per acre to control thrips. Do not apply within 10 days of harvest.

POTATOES: Apply 20-50 lbs. per acre to control aphids, flea beetles, Colorado potato beetles, dipterous leaf miners, banded cucumber beetles, leafhoppers, southern armyworms. Do not apply within 14 days of harvest. Foliage application.

TOMATOES: Apply 20-40 lbs. per acre to control aphids, dipterous leaf miners, banded cucumber beetles, fall armyworms, southern armyworms, vinegar fly. Do not apply within 3 days of harvest.

ALFALFA, CLOVER: Apply 15 lbs. per acre to control spotted alfalfa aphids, yellow clover aphids, pea aphids, leafhoppers, plant bug nymphs, mites. Do not apply within 7 days of cutting or grazing by animals being finished for slaughter.

CORN: Apply 30-40 lbs. per acre to control grasshoppers, corn leaf aphid, corn earworms. (Apply when silks first appear. Two or three applications at two or three day intervals will be necessary.) Do not apply within 2 days of harvest. Do not feed forage to livestock within 10 days following last application.

COTTON: Apply 20-25 lbs. per acre to control spider mites, aphids, leafhoppers, cotton leaf worms, cotton leaf perforators, lygus, salt marsh caterpillars. Do not apply within 14 days of harvest. Do not graze dairy or meat animals in treated fields. Do not feed gin waste to livestock.

PUREGRO CONDITIONS OF SALE AND WARRANTIES*

PureGro Company warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label. PureGro makes no other warranty, express or implied, of fitness or merchantability nor authorizes any agent or representative to make any variation or exception from this warranty.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of PureGro Company. Buyer assumes all risk of use, storage, or handling of this material not in strict accordance with directions given herewith.

ORM-A

**RQ DIAZINON
MIXTURE NA2783**

Reg # 1202-152
PM 15
1-20-72