

ACCEPTED

MAY 5 - 1967

1202-153

DIRECTIONS FOR USE

Apply when air is calm, using ground equipment or aircraft. Make thorough coverage of plants with a uniform deposit of dust. Apply at the first sign of infestation and repeat at regular intervals when needed to maintain control, but observe use limits given for specific crops.

Fruit and Nut Crops

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES: Apply 15-100 lbs. per acre to control mites, aphids. Do not apply within 1 day of harvest.

FIGS: Apply 50 lbs. per acre to control dried fruit beetles, vinegar fly. Do not apply within 3 days of harvest.

FILBERTS: Apply 50-75 lbs. per acre to control filbert aphid, apple mealybug, eye-spotted bud moth. No time limitation.

GRAPES: Apply 30-60 lbs. per acre to control *Drosophila*, leafhopper. Do not apply within 3 days of harvest.

PECANS: Apply 100 lbs. per acre to control pecan nut case bearer, pecan bud moth. No time limitation.

STRAWBERRIES: Apply 25-50 lbs. per acre to control lygus bug, mille bug, leaf crickets, thrips, potato leafhopper, strawberry leaf roller, white fly. Apply 40-50 lbs. per acre to control aphids, spider mites. Do not apply within 3 days of harvest.

WALNUTS: Apply 60-100 lbs. per acre to control aphid. No time limitation.

Field, Forage & Vegetable Crops

ALFALFA: Apply 30-35 lbs. per acre to control aphid, potato leafhopper, spider mites, alfalfa weevil larvae, spittle bug, vetch bruchid, grasshopper. Apply 15-30 lbs. per acre to control spotted alfalfa aphid. No time limitation.

BARLEY, OATS, RYE, WHEAT: Apply 10 lbs. per acre to control English grain aphid, winter grain mite, young grasshoppers. Do not apply within 7 days of harvest.

BEANS: Apply 30-40 lbs. per acre to control Mexican bean beetle, leafhoppers, spider mites, Japanese beetle, aphids, cucumber beetles. Do not apply within 1 day of harvest.

CARROTS: Apply 35-50 lbs. per acre to control aphids, leafhoppers. Do not apply within 7 days of harvest.

CELERY: Apply 35 lbs. per acre to control aphids, spider mites. Do not apply within 7 days of harvest.

CLOVER: Apply 25-45 lbs. per acre to control aphids, mites, young grasshoppers, leafhopper, clover leaf weevil. No time limitation.

CRUCIFERS (Broccoli, Brussels sprouts, Cabbage, Cauliflower): Apply 30-60 lbs. per acre to control aphids, beetles, imported cabbage worm, cabbage looper, diamondback. Do not apply within 7 days of harvest, 3 days of harvest on broccoli.

COLLARD: Apply 30-60 lbs. per acre to control aphid, harlequin cabbage bug, imported cabbage looper, cabbage looper, diamondback moth. Do not apply within 7 days of harvest.

CORN (Field, Pop, Sweet): Apply 40 lbs. per acre to control sap beetle, corn earworm, thrip, aphid, young grasshopper. Do not apply within 5 days of harvest. May cause injury in the whorl stage.

(Continued on right panel)



MALATHION 4 DUST

ACTIVE INGREDIENT: Malathion (0, 0-dimethyl dithiophosphate or diethyl mercaptosuccinate) 4.0%

INERT INGREDIENTS: 96.0%

100.0%

CAUTION

AVOID BREATHING DUST.

AVOID REPEATED CONTACT WITH SKIN.

WASH THOROUGHLY AFTER USING.

AVOID CONTAMINATION OF FEED AND FOODSTUFFS.

OBSERVE FOLLOWING USE PRECAUTIONS

Observe use limitations given on the label for specific crops. Do NOT apply during bloom when honey bees or other pollinators are active. Avoid dust drift to adjoining food and forage crops about to be harvested, or crops in bloom. Apply in early morning or evening when there is little wind. The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified.

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label. Containers should be disposed of by burying with wastes, or by burning (keep out of smoke).

KEEP OUT OF REACH OF CHILDREN.



Net Weight _____

(Continued from left panel)

CITRUS: Apply 15-100 lbs. per acre to control spider mite, cotton aphid, leafhopper, white fly, cotton leafworm, brown cotton leafworm, cotton leaf perforator, thrips, lygus bug, flea hopper. Add 10 lbs. actual DDE to control bollworm. No time limitation.

EGG PLANT: Apply 30-50 lbs. per acre to control aphids, lace bug. Do not apply within 3 days of harvest.

ENJOE: Apply 30-60 lbs. per acre to control aphid, spider mite. Do not apply within 7 days of harvest.

LETTUCE: Apply 30-100 lbs. per acre to control aphid, mites, cabbage looper, leafhopper. Do not apply within 14 days of harvest or leaf fall, or 7 days of harvest on leaf lettuce.

ONIONS: Apply 30-50 lbs. per acre to control onion thrips. Do not apply within 14 days of harvest on green onions.

PEANUTS: Apply 25 lbs. per acre to control potato leafhopper. Do not apply within 30 days of harvest.

PEAS: Apply 25-60 lbs. per acre to control pea aphid, pea weevil. Do not apply within 7 days of harvest if vines are to be fed to livestock. 14 days of harvest if vines are not for forage.

PEPPERS: Apply 15 lbs. per acre to control aphids, pepper maggot. Do not apply within 14 days of harvest.

POTATOES: Apply 25-75 lbs. per acre to control aphid, leafhopper, potato starch bug. Use 30-75 lbs. per acre to control weevils. No time limitation.

PUMPKINS: Apply 30-50 lbs. per acre to control aphid, mites, spider mites. Use 50 lbs. per acre to control squash vine borer. Do not apply within 1 day of harvest. Do not apply to young plants or fruit.

RADISHES, RABBITS, SALADY, CUCUMBERS, RADISHES: Apply 30-50 lbs. per acre to control aphids. Do not apply within 7 days of harvest.

SCALOPES: Apply 30-60 lbs. per acre to control aphids. Do not apply within 14 days of harvest.

SQUASH: Apply 30-50 lbs. per acre to control squash vine borer, aphid, spider mites, weevil, worm, flea hopper. Do not apply to control squash vine borer on fruit or stems. Do not apply within 7 days of harvest. Do not apply to young plants or fruit.

SPINACH: Apply 30-50 lbs. per acre to control aphids. Do not apply within 7 days of harvest.

SUGAR BEETS: Apply 30-50 lbs. per acre to control aphid, grasshopper, grasshopper. Do not apply within 7 days of harvest if top parts to be used for food.

SWISS CHARD: Apply 30-50 lbs. per acre to control aphid, grasshopper, grasshopper. Do not apply within 7 days of harvest.

VEGETABLES: Apply 30-50 lbs. per acre to control grasshopper, grasshopper, grasshopper, grasshopper. Do not apply within 7 days of harvest.

WATERMELONS: Apply 30-50 lbs. per acre to control grasshopper, grasshopper, grasshopper, grasshopper. Do not apply within 7 days of harvest.

WHEAT: Apply 10 lbs. per acre to control English grain aphid, winter grain mite, young grasshoppers. Do not apply within 7 days of harvest.

YARROW: Apply 30-50 lbs. per acre to control aphid, grasshopper, grasshopper, grasshopper. Do not apply within 7 days of harvest.