[illegible]

[illegible] per acre to control mites, peach [illegible] apply after buds open. Do not feed hulls [illegible]

[illegible] per acre to control mites, aphids, codling [illegible] fruit tree leaf roller, [illegible] days of harvest.

[illegible] Apply 25-10 [illegible] apply within 15 days of harvest.

[illegible] per acre to control mites, aphids, [illegible] fruit worms, oriental [illegible] days of harvest.

[illegible] lemons, tangerines, [illegible] control mites, citrus [illegible] Apply

[illegible] per acre to control [illegible] fruit tree leaf roller, [illegible] plum curculio. [illegible] Do not [illegible] apply within [illegible] actual [illegible] Do not apply within 21 days of harvest. Do not apply [illegible] Do not apply [illegible] Do not graze treated [illegible] land.

[illegible] per acre to control mites, pear psylla, [illegible]

[illegible] per acre to control mites, [illegible] heavy lime sulfur. Leafhopper, [illegible] days of harvest. Do not [illegible]

[illegible] per acre to control mites [illegible] within 21 days of harvest.

[illegible] per acre to control mites, thrips, [illegible] within 21 days of harvest.

(Continued on next panel)

Repackaged

Manufactured by
PACIFIC GUANO COMPANY

KELTHANE PARATHION
3-2 DUST

ACTIVE INGREDIENTS:

1,1'-bis (p-chlorophenyl)-2,2,2-trichloroethanol	3.0%
Parathion (O, O-diethyl O-p-nitrophenyl thiophosphate)	2.0%
INERT INGREDIENTS:	96.0%
	100.0%

® U.S. Reg. U.S. Pat. Off. by Rohm & Haas Company

WARNING

KEEP OUT OF REACH OF CHILDREN.

May be Fatal If Swallowed, Inhaled, or Absorbed Through Skin. Rapidly Absorbed Through Skin. Do not get in eyes, on skin, or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Treated areas should not be re-entered until drilling insecticide and volatile residues have dissipated. Do not contaminate feed and feedstuffs. Wash hands, arms, and face thoroughly with soap and water before eating, or smoking. Wash all contaminated clothing with soap and hot water before re-use.

This product is toxic to fish and wildlife.

Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label.

Manufactured by
PACIFIC GUANO COMPANY

BURN EMPTY CONTAINER
STAY OUT OF SMOKE

Net Contents [illegible]

ACCEPTED
MAY 9 1967
FOR THE ADMINISTRATOR
[illegible]
ENVIRONMENTAL PROTECTION AGENCY
REGISTRATION NO. 432-116
SEE ATTACHED COMMENTS

(Continued from left panel)

CORN (Field corn only): Apply 2 to 50 lbs. per acre to control aphid, mites, armyworm, budworm, European corn borer, grasshoppers, sap beetle. Do not apply after ears begin to form. Do not feed treated corn or use it as forage.

COTTON: Apply 2½ to 40 lbs. per acre to control mites, aphids, leafhoppers, garden webworm. Do not apply within 14 days of harvest. Do not graze treated fields by meat or dairy animals.

CUCUMBERS: Apply 25 lbs. per acre to control mites, cucumber beetle, melonworm, pickleworm, serpentine leaf miner. Do not apply within 15 days of harvest. Do not apply before vines. Do not apply to cucurbits unless plants are dry.

EGGPLANTS: Apply 20 to 30 lbs. per acre to control mites, aphid, leaf miner. Do not apply within 14 days of harvest.

HOPS: Apply 2 to 40 lbs. per acre to control mites, aphids. Do not apply within 14 days of harvest.

MELONS, PUMPKINS, WATERMELONS: Apply 25 lbs. per acre to control aphids, mites, stink bug, vine borer, cucumber beetle, melonworm, pickleworm, serpentine leaf miners. Do not apply within 10 days of harvest. 7 days of harvest melons.

PEPPERS: Apply 2 to 30 lbs. per acre to control aphid, mites, serpentine leaf miner. Do not apply within 14 days of harvest.

POTATOES: Apply 2 to 40 lbs. per acre to control mites. Colorado potato beetle, flea beetle, leafhopper, serpentine leaf miner, whitefly, grasshoppers, aphid, armyworms. Do not apply within 7 days of harvest.

SQUASH (Summer and Winter): Apply 2 lbs. per acre to control mites, cucumber beetle, melonworm, pickleworm, aphid, serpentine leaf miner. Do not apply within 14 days of harvest.

STAKE BEANS: Apply 2 to 30 lbs. per acre to control mites, aphid, web worm. Do not apply within 21 days of harvest. Do not feed vines to dairy animals or meat animals.

TOMATOES: Apply 25 to 30 lbs. per acre to control Colorado potato beetle, flea beetle, leafhopper, serpentine leaf miner, grasshoppers, whitefly, aphids, armyworms, mites. Do not apply within 14 days of harvest.

NONWARRANTY

Because the use of this material is subject to conditions beyond Pacific Guano Company's control, neither Pacific Guano Company, nor any other Seller hereof makes any warranty, express or implied, of use thereof or the results of such use when material is used in accordance with directions. No Seller is responsible for the results of any use of this material.

USDA Reg. No. [illegible]