

...on the amount of 2,4-dimethyl para-  
...out water to cover the entire surface...

...to control adults...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...

...METHYL PARATHION...



# METHYL PARATHION

## 25 WP

### ACTIVE INGREDIENTS

\*0, 0-dimethyl 0-p-nitrophenoxyl phosphorothioate 25.00%

### INERT INGREDIENTS

\*Methyl Parathion

### KEEP OUT OF REACH OF CHILDREN.

**POISONOUS BY SKIN CONTACT, INHALATION OR SWALLOWING  
RAPIDLY ABSORBED THROUGH SKIN  
REPEATED EXPOSURE MAY, WITHOUT SYMPTOMATIC  
E INCREASINGLY HAZARDOUS**

**Do not get in eyes, on skin, on clothing.**

Wear clean, heavy, natural rubber gloves and clean, waterproof or freshly laundered, protective clothing (overall, cap, etc.) and goggles. Destroy and replace gloves frequently. Wash hands, face and arms thoroughly with soap and warm water before eating or smoking. If the immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before re-use.


In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water.

**Do not breathe dust, vapor or spray mist.**

Wear mask of respirator type passed by the U.S. Department of Agriculture for methyl parathion use. For protection, use any one of the following: full-face canister type mask; dust-proof hood with mechanical filter; dust ventilation.

**Do not store near food or feed stuffs.**

Apply in place of recommended rate with all anti-solvent and then with water. Do not apply to any surface that is out of sight. Do not use in case of all unauthorized persons, all of general area or vicinity where there may be stay of children. Do not apply to any surface that is out of sight without a suitable protective covering. Do not use in case of any other person, child, animal, or plant, or any other surface that is out of sight without a suitable protective covering.



PACIFIC CYANIDE COMPANY  
100-108-100-100

... ..

### DANGER

(Continued from middle panel)

### WARNING

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label.

Do not apply during bloom when honey bees or other pollinators are active. Avoid drift to adjoining fields, gardens, woods or other areas.

### FIRST AID TREATMENT

If symptoms or signs of poisoning include blurred vision, difficulty breathing, chest tightness, do not wait for a doctor. Give 2 atropine tablets (each 1/100 gr.) at once. Remove patient immediately from the area where methyl parathion is present. Remove contaminated clothing and wash the soiled areas with plenty of soap and water to remove all traces of parathion. If parathion gets in eyes, immediately flush with water for at least 15 minutes and get medical attention. If swallowed, give 1-2 spoonfuls of milk with water and repeat until vomit fluid is clear. Have victim lie down and keep quiet.

**NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.** (See the special Note.) Warning symptoms include weakness, headache, tightness in the chest, blurred vision, non-reactive pupils, rapid salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

**TREATMENT:** Give atropine, preferably by injection, grains 1-100, 2-30. Tablets at once and parenterally or orally every hour until a normal pulse is obtained or until pupils dilated. Never give morphine. If at all else by postural drainage. Artificial respiration or oxygenation treatment may be necessary. Observe patient continuously for 48 hours. Repeat exposure to cholinesterase inhibitor may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for a cholinesterase re-examination has been allowed to determine the blood level.

### NOTICE

Because this product is a pesticide, it is subject to conditions of sale. The product is sold by the Pacific Cyanide Company, a subsidiary of the Cyanide and Chemicals Corporation, and is sold only to those who are qualified to use it. The results of such use, whether or not used in accordance with the directions, are not warranted, with or without a contract. Neither of us is responsible for any loss or damage resulting from the use of this product.

USA 100-108-100-100