

DIRECTIONS FOR USE

Field Crops and Vegetables

COTTON: Thrips, flea beetle, fleahopper, leafworm, striped blister beetle, boll weevil, bollworm, fall armyworm, leaf perforator, leaf rollers, leafhoppers, tarnished plant bug, light to moderate numbers of Western Lygus bugs, aphids will be repelled by repeat applications in a schedule), pink bollworm, salt marsh caterpillar, stink bugs, grasshoppers--Apply 20-23 lbs. per acre. No time limitations. Allow 7 days before grazing or harvesting crop residue for feed or bedding for dairy or meat animals.

BEANS: Mexican bean beetle, leafhoppers, bean leaf beetle, cucumber beetles, Western bean cutworm, flea beetles, Japanese beetle, velvet-bean caterpillar, corn earworm, armyworms, stink bugs, tarnished plant bug, (lima bean pod borer) corn earworm, Lygus and stink bugs--Apply 15-20 lbs. per acre. No time limitation.

PEAS: Leafhoppers, Colorado potato beetle, armyworms, grasshoppers--Apply 15 lbs. per acre. No time limitation.

CORN: Corn earworm, (Apply to silks as soon as they appear. Make 3 to 4 application at 2-3 day intervals. Re-apply immediately after rain.), corn rootworm adults, European corn borer, fall armyworm, sap beetles, Japanese beetle, flea beetles, leafhoppers--Apply 15-20 lbs. per acre. No time limitation.

SOYABEANS: Leafhoppers, armyworms, webworms, grasshoppers--Apply 15-20 lbs. per acre. Do not apply 14 days before harvest.

POTATO: Colorado potato beetle, flea beetles, leafhoppers--Apply 15-20 lbs. per acre. Foliage application only.

TOMATO, EGGPLANT, PEPPER: Tomato fruitworm, fall armyworm, tomato hornworm, European corn borer, stink bugs, lace bugs, Lygus bugs, Colorado potato beetle, flea beetles, leafhoppers--Apply 15-20 lbs. per acre. No time limitation.

ASPARAGUS: Apply 15-20 lbs. per acre to control asparagus beetle. Apply during cutting season. Do not repeat application within 3 days. Do not apply 1 day before harvest.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI: Flea beetle, harlequin bug, corn earworm, armyworms. For small loopers after heads form, use SEVIN every 5 to 7 days--Apply 15-20 lbs. per acre. Do not apply 3 days before harvest.

Lot No. _____



**SEVIN SULFUR
DUST 10-25**

ACTIVE INGREDIENTS:	
Carbaryl (1-naphthyl-N-methylcarbamate)	10.0%
Sulfur	25.0%
INERT INGREDIENTS:	65.0%
KEEP OUT OF REACH OF CHILDREN.	100.0%

CAUTION

Harmful if swallowed. Avoid breathing dust. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid storage near feed and food products.

This product is toxic to fish. Keep out of lakes, ponds and streams.

Do not apply to any area not specified on the label.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.



Registered trademark of Union Carbide Corporation for the active ingredient (Carbaryl)

BURN EMPTY CONTAINER
KEEP OUT OF SMOKE

NET CONTENTS _____

Tree and Shrub Crops

APPLES, PEARS: Codling moth, green apple aphid, pear psylla, orange tortrix, California pear slug. Apply 15-100 lbs. per acre. Do not apply 1 day before harvest.

PEACHES, APRICOTS, NECTARINES: Oriental fruit moth, peach twig borer, codling moth, orange tortrix, tussock moth, fruit tree leaf roller, peritrocha cicada, Japanese beetle. Apply 40-80 lbs. per acre. Do not apply 1 day before harvest for peaches and 3 days before harvest for apricots and nectarines.

GRAPES: Grape leaf folder, grape leafhopper, grape leaf skeletonizer, grape berry moth, Japanese beetle, June beetles, red-banded leaf roller--Apply 20-30 lbs. per acre. No time limitation.

CHERRY, PLUM, PRUNE: Cherry fruit fly, fruit tree leaf roller, peach twig borer, orange tortrix, Western tussock moth--Apply 40-60 lbs. per acre. Do not apply 1 day before harvest.

FILBERT: Filbert worm, filbert aphid, filbert leaf roller--Apply 50 lbs. per acre. No time limitation.

Forage and Feed Crops

ALFALFA, CLOVERS, PEANUTS, SOYBEANS, COWPEAS: Blister beetles, Mexican bean beetle, thrips, leafhoppers, three-cornered alfalfa hopper, cucumber beetles, bean leaf beetle, Japanese beetle, alfalfa caterpillar, velvetbean caterpillar and green cloverworm, armyworms, webworms, corn earworm, stink bugs, grasshoppers. Apply 15 lbs. per acre. No time limitation.

SORGHUM (Grain and Forage): Armyworm, webworm, corn earworm, stink bugs, sorghum midge, grasshoppers--Apply 20 lbs. per acre. Do not apply 21 days before harvest of grain. No time limitation on sorghum forage.

RICE, FORAGE GRASSES, JUGLANS: Armyworms, webworms, thrips. Apply 15 lbs. per acre. Do not allow thrip control to progress to row for seed, dust into holes. Do not apply to Rice within 10 days of harvest. May be applied up to 10 days after harvest of rice.

NONWARRANTING

Because the use of this material is subject to conditions beyond the control of the Pacific Guard Company, no warranty, either expressed or implied, is made by the use thereof or the results of such use, whether or not used in accordance with the directions. Neither of the foregoing will or shall be responsible for the results of any use of this material.

USDA Reg. No. _____

14-2-162
MAY 11 1967

ACCEPTED

MAY 11 1967

IN ACCORDANCE WITH FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTICIDE REGISTERED
UNDER NO. 202-165 SUBJECT
TO ATTACHED COMMENTS

120-165

DIRECTIONS FOR USE



**SEVIN SULFUR
DUST 10-75**

ACTIVE INGREDIENTS:	
Carbaryl (1-naphthyl-N-methylcarbamate)	10.0%
Sulfur	75.0%
INERT INGREDIENTS	15.0%

KEEP OUT OF REACH OF CHILDREN. 100.0%

CAUTION

Harmful if swallowed. Avoid breathing dust. Avoid prolonged or repeated contact with skin. Wash thoroughly after using. Avoid storage near feed and food products.

This product is toxic to fish. Keep out of lakes, ponds and streams.

Do not apply to any area not specified on the label.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.



Registered trademark of Union Carbide Corporation for the active ingredient (Carbaryl).

BURN EMPTY CONTAINER
KEEP OUT OF SMOKE

Field Crops and Vegetables

COTTON: Thrips, flea beetle, leafhopper, leafworm, striped blister beetle, boll weevil, bollworm, fall armyworm, leaf perforator, leaf rollers, leafhoppers, tarnished plant bug, light to moderate numbers of Western Lygus bug. Aphids will be repressed by repeat applications in a schedule; pink bollworm, salt marsh caterpillar, stink bugs, grasshoppers. Apply 20-23 lbs. per acre. No time limitations. Allow 7 days before grazing or harvesting crop residue for feed or bedding for dairy or meat animals.

BEANS: Mexican bean beetle, leafhoppers, bean leaf beetle, cucumber beetles, Western bean cutworm, flea beetles, Japanese beetle, velvet-bean caterpillar, corn earworm, armyworms, stink bugs, tarnished plant bug, lima bean pod borer, corn earworm, Lygus and stink bugs. Apply 15-20 lbs. per acre. No time limitation.

PEAS: Leafhoppers, Colorado potato beetle, armyworms, grasshoppers. Apply 15 lbs. per acre. No time limitation.

CORN: Corn earworm, (Apply to silks as soon as they appear. Make 3 to 4 application at 2-3 day intervals. Re-apply immediately after rain.), corn rootworm adults, European corn borer, fall armyworm, sap beetles, Japanese beetle, flea beetles, leafhoppers. Apply 15-20 lbs. per acre. No time limitation.

SUGAR BEETS: Leafhoppers, armyworms, webworms, grasshoppers. Apply 15-20 lbs. per acre. Do not apply 14 days before harvest.

POTATO: Colorado potato beetle, flea beetles, leafhoppers. Apply 15-20 lbs. per acre. Forage application only.

TOMATO: European corn borer, fall armyworm, tomato hornworm, European corn borer, stink bugs, lace bugs, Lygus bugs, Colorado potato beetle, flea beetles, leafhoppers. Apply 15-20 lbs. per acre. No time limitation.

ASPARAGUS: Apply 15-20 lbs. per acre to control asparagus beetle. Apply during cutting season. Do not repeat application within 3 days. Do not apply 1 day before harvest.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABE: Flea beetle, harlequin bug, corn earworm, armyworms. For small loopers after heads form, use SEVIN every 5 to 7 days. Apply 15-20 lbs. per acre. Do not apply 3 days before harvest.

Lot No. _____

NET CONTENTS _____

FRUIT AND NUTS

APPLES, PEARS: Codling moth, green apple aphid, pear psylla, orange tortrix, California pear slug. Apply 15-20 lbs. per acre. Do not apply 1 day before harvest.

PLUMS, APRICOTS, NECTARINES: Oriental fruit moth, peach twig borer, codling moth, orange tortrix, tussock moth, fruit tree leaf roller, periwinkle weevil, Japanese beetle. Apply 40-50 lbs. per acre. Do not apply 1 day before harvest for peaches and 3 days before harvest for apricots and nectarines.

GRAPES: Grape leaf folder, grape leafhopper, grape leaf skeletonizer, grape berry moth, Japanese beetle, June beetles, red-banded leaf roller. Apply 20-30 lbs. per acre. No time limitation.

CHERRY, PLUM, PRUNE: Cherry fruit fly, fruit tree leaf roller, peach twig borer, orange tortrix, Western tussock moth. Apply 40-50 lbs. per acre. Do not apply 1 day before harvest.

FILBERT: Filbert worm, filbert aphid, filbert leaf roller. Apply 50 lbs. per acre. No time limitation.

Forage and Feed Crops

ALFALFA, CLOVERS, PEANUTS, SOYBEANS, COWPEAS: Blister beetle, Mexican bean beetle, thrips, leafhopper, three-cornered alfalfa hopper, cucumber beetles, bean leaf beetle, Japanese beetle, alfalfa caterpillar, cowpea caterpillar and green cloverworm, armyworms, webworms, corn earworm, stink bugs, grasshoppers. Apply 15 lbs. per acre. No time limitation.

SORGHUM (Grain and Forage): Armyworm, webworms, corn earworm, stink bugs, sorghum midge, grasshoppers. Apply 20 lbs. per acre. Do not apply 2 days before harvest of grain. No time limitation on sorghum forage.

RICE, FORAGE GRASSES, PASTURE: Armyworms, stink bugs, etc. Apply 15 lbs. per acre. For optimum thrip control on grasses, apply for seed. Apply just into boot. Do not apply to Rice within 14 days of harvest. May be applied up to harvest on Pastures and Forage Grasses.

NO WARRANTY

Because the use of this material is subject to conditions beyond their control, neither Pacific Guard Company, nor any other Seller hereon, warrant or other express or implied the use thereof or the results of such use, whether or not use is in accordance with the directions. Neither of the foregoing will or shall be responsible for the results of any use of this material.

USDA Reg. No. _____