DIRECTIONS FOR USE

Apply when air is calm, using ground equipment or aircraft. Make thorough coverage of plants with a uniform deposit of dust. Apply at the first sign of infestation and repeat at regular intervals when needed to maintain control, but observe use limits given for specific crops.

Fruit and Nut Crops

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES: Apply 35-40 lbs. per acre to control mites, aphids. Do not apply within 1 day of harvest.
FIGS: Apply 40 lbs. per acre to control dried fruit beetles, vinegar fly. Do not apply within 3 days of harvest.
FILBERTS: Apply 40-60 lbs. per acre to control filbert aphid, apple mealybug, eye-spotted bud moth. No time limitation.
GRAPES: Apply 25-50 lbs. per acre to control Drosophila, leafhopper. Do not apply within 3 days of harvest.
PECANS: Apply 200 lbs. per acre to control pecan nut case bearer, pecan bud moth. No time limitation.
STRAWBERRIES: Apply 20-40 lbs. per acre to control lygus bug, spittle bug, field crickets, thrips, potato leafhopper, strawberry leaf roller, white fly. Apply 30-40 lbs. per acre to control aphids, spider mites. Do not apply within 3 days of harvest.
WALNUTS: Apply 50-200 lbs. per acre to control aphid. No time limitat[illegible].

Field, Forage & Vegetable Crops

ALFALFA: Apply 25-40 lbs. per acre to control aphid, potato leafhopper, spider mites, alfalfa weevil larvae, spittle bug, vetch bruchid, grasshopper. Apply 15-30 lbs. per acre to control spotted alfalfa aphid. No time limitation.
BARLEY, OATS, RYE, WHEAT: Apply 25 lbs. per acre to control English grain aphid, winter grain mite, young grasshoppers. Do not apply within 7 days of harvest.
BEANS: Apply 25-35 lbs. per acre to control Mexican bean beetle, leafhoppers, spider mites, Japanese beetle, aphids, cucumber beetles. Do not apply within 1 day of harvest.
CARROTS: Apply 30-40 lbs. per acre to control aphids, leafhoppers. Do not apply within 7 days of harvest.
CELERY: Apply 30 lbs. per acre to control aphids, spider mites. Do not apply within 7 days of harvest.
CLOVER: Apply 20-30 lbs. per acre to control aphids, mites, young grasshoppers, leafhopper, clover leaf weevil. No time limitation.
CRUCIFER CROPS (Broccoli, Brussels Sprouts, Cabbage, Cauliflower): Apply 25-50 lbs. per acre to control aphids, mites, imported cabbage worm, cabbage looper, diamondback. Do not apply within 7 days of harvest, 3 days of harvest on broccoli.
COLLARD: Apply 25-50 lbs. per acre to control aphid, harlequin cabbage bug, imported cabbage worm, cabbage looper, diamondback moth. Do not apply within 7 days of harvest.
CORN (Field, Pop, Sweet): Apply 30 lbs. per acre to control sap beetle, corn earworm, corn aphid, young grasshopper. Do not apply within 5 days of harvest. May cause injury in the whorl stage.
COTTON: Apply 15-30 lbs. per acre for control of spider mites, cotton aphid, leafhoppers, white fly, cotton leafworm, brown cotton leafworm, cotton leaf perforator, thrips, lygus bugs, flea hoppers. No time limitation.
CUCUMBERS, MELONS (Cantaloupes, Casabas, Crenshaw, Honeydew, Honeyballs, Musk melon, Persian, Watermelon): Apply 25-40 lbs. per acre to control aphid, spider mites, pickleworm. Apply 35-40 lbs. per acre to control cucumber beetles. Do not apply within 1 day of harvest. Do not apply to cucurbits unless plants are dry.
EGGPLANT: Apply 10-70 lbs. per acre to control aphids, lace bug. Do not apply within 3 days of harvest.

Lot No. _____ 24067R/W

GAVICIDE

MALATHION 5 DUST

ACTIVE INGREDIENT:	
Malathion (O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate)	5.0%
INERT INGREDIENTS:	95.0%
	100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN.

AVOID BREATHING DUST.
AVOID REPEATED CONTACT WITH SKIN.
WASH THOROUGHLY AFTER USING.
AVOID CONTAMINATION OF FEED AND FOODSTUFFS.

OBSERVE FOLLOWING USE PRECAUTIONS

Observe use limitations given on the label for specific crops.
Do NOT apply during bloom when honey bees or other pollinators are active.
Avoid dust drift to adjoining food and forage crops
Apply in early morning or evening when there is little wind.
The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified.

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label. Containers should be disposed by burying with wastes, or by burning (Keep out of smoke).

Manufactured By
PUREGRO
LOS ANGELES, CALIFORNIA

Net Weight 50 _____ lbs.

ACCEPTED
12/8/67
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 1202-164 SUBJECT TO ATTACHED COMMENTS.

(Continued From Left Panel)

ENDIVE: Apply 25-40 lbs. per acre to control aphid, spider mite. Do not apply within 7 days of harvest.
LETTUCE: Apply 25-80 lbs. per acre to control aphid, mites, cabbage looper, leafhoppers. Do not apply within 14 days of harvest on leaf lettuce, 7 days of harvest on head lettuce.
ONIONS: Apply 25-40 lbs. per acre to control onion thrip. Do not apply within 3 days of harvest on green onions.
PEANUTS: Apply 20 lbs. per acre to control potato leafhopper. Do not apply within 30 days of harvest.
PEAS: Apply 20-50 lbs. per acre to control pea aphids, pea weevil. Do not apply within 7 days of harvest if vines are to be fed to livestock, 3 days of harvest if vines are not for forage.
PEPPERS: Apply 30 lbs. per acre to control aphids, pepper maggots. Do not apply within 3 days of harvest.
POTATOES: Apply 20-60 lbs. per acre to control aphid, leafhopper, false shinch bug. Use 50-75 lbs. per acre to control mealybug. No time limitation.
PUMPKINS: Apply 30-40 lbs. per acre to control aphids, mites, leafhoppers. 40 lbs. per acre to control squash vine borer. Do not apply within 3 days of harvest. Do not apply unless plants are dry.
RADISHES, PARSNIPS, SALSIFY, HORSERADISHES: Apply 25-40 lbs. per acre to control aphids. Do not apply within 7 days of harvest.
RUTABAGAS: Apply 25-50 lbs. per acre to control aphids. Do not apply within 3 days of harvest.
SOYBEANS: Apply 30-40 lbs. per acre to control Mexican bean beetle. Do not apply within 3 days of harvest.
SQUASH: Apply 30-40 lbs. per acre to control alfalfa loopers, aphid, spider mites, pickle worm, 35-40 lbs. per acre to control squash vine borer, cucumber beetle. Do not apply within 7 days of harvest. Do not apply unless plants are dry.
SPINACH: Apply 25-40 lbs. per acre to control aphids. Do not apply within 7 days of harvest.
SUGAR BEETS: Apply 25-50 lbs. per acre to control aphids, spider mites, apply 40-50 lbs. to control grasshoppers. Do not apply within 7 days of harvest if tops are to be used for food or feed.
SWEET POTATOES: Apply 25-40 lbs. per acre to control leafhopper. No time limitation.
TOMATOES: Apply 30-40 lbs. per acre for control of spider mites, aphids, tomato russet mite, Drosophila. Do not apply within 3 days of harvest.
TURNIP, KALE: Apply 25-50 lbs. per acre to control aphids, imported cabbage worm, cabbage looper. Do not apply within 7 days of harvest on kale, 3 days of harvest on turnips.
FLOWERS (Stock, Iris, Daisies, Mums, Gladiolus, Astera): Apply 20-50 lbs. per acre to control spider mites, white fly, mealybugs.

[illegible] Manufacturer warrants that this material conforms to the chemical description on the label and is reasonably suitable for the purposes stated on the label when used in accordance with the directions under normal conditions of use. Manufacturer makes no other warranty of FITNESS OR MERCHANTABILITY, guarantee or representation expressed or implied, concerning this material. In case of material defect or failure to conform to the manufacturer's guarantee, buyer's exclusive remedy shall be limited to the return of the material or, at the option of the manufacturer, the refund of the purchase price. In no case shall manufacturer be liable for consequential or special damages resulting from the use or handling of this material. Buyer and user accept this material subject to the foregoing conditions of sale which cannot be changed except by agreement in writing signed by the manufacturer's authorized agent. Buyer and user assume all risks and liability whatsoever for the use, storage or handling of this material not in accordance with the directions or warnings, even though such use, storage or handling may be the sole cause of loss, damage or injury. Buyer and user agree to indemnify manufacturer against all claims and liability arising from handling, storage and use of this material not in accordance with such directions or warnings. Buyer and user assume all risks and liability whatsoever (except as to claims for personal injury) resulting from handling, storage and use of this material not in accordance with such directions or warnings, even though such handling, storage or use may be the sole cause of loss, damage or injury. Buyer and user agree to indemnify manufacturer against all claims and liability arising from handling, storage and use of this material not in accordance with such directions or warnings.