

DIRECTIONS FOR USE

APPLES, PEARS: Apply 85-200 lbs. per acre to control powdery mildew and to suppress European red mite, clover and two spotted mite. Begin application at delayed dormant to pink stage and continue 7 to 14 day intervals until terminal growth or mildew activity is completed. Do not apply within 21 days of harvest. Do not apply within 2 to 3 weeks of an oil spray and do not use on sulfur sensitive varieties.

PEACHES: Apply 75-100 lbs. per acre to control powdery mildew. Can be used at pink bud, petal fall, shuck fall and early covers. Do not apply within 45 days of harvest.

CANTALOUPE, CUCUMBERS, HONEY-DEW MELONS, MUSMELONS, PUMPKINS, SQUASH, WATERMELONS: Apply 40-50 lbs. per acre to control powdery mildew. Begin application when mildew first appears and repeat at 7 to 10 day intervals as long as disease threatens. On summer squash and cucumber -- a maximum of 4 applications should be applied to an individual set of fruit. Do not apply within 7 days of harvest.

GRAPES: Apply 50-75 lbs. per acre to control powdery mildew. Repeat at 10 to 14 day intervals as long as disease threatens. Do not use on Thompson seedless or Pearl varieties. Do not apply within 21 days of harvest.

TREES, ORNAMENTAL TREES AND SHRUBS, FLOWERS: Apply 40-75 lbs. per acre to control powdery mildew, red spider mite, clover mite.

Lot No. _____ R24467RW



KARATHANE

SULFUR DUST 1-30

ACTIVE INGREDIENTS:

Dinitro (1-methylheptyl) phenyl <u>cronate</u>	0.9%
Other nitro phenols and derivatives, chiefly dinitro (1-methylheptyl)* phenol	0.1%
Sulfur	50.0%
INERT INGREDIENTS	49.0%

100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN.

May cause irritation. Avoid contact with skin and eyes. Avoid inhalation of dust. In case of contact, wash thoroughly. Do not take internally. Avoid contamination of foodstuffs. Wash or wipe edible portions of fruits or vegetables thoroughly before use. Do not use when temperatures exceed 90° F. Apply this product only as specified on this label.

(Continued on right panel)



NET CONTENTS 50 lbs.

(Continued from front panel)

CAUTION

This product will kill fish. Keep out of any body of water.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by burying with wastes, or by burning (Keep out of smoke).

Do not apply to cucurbits unless the variety is known to be sulfur tolerant under local conditions. Avoid drift to sulfur sensitive plants.

1. PURCHASE OF SALES: Puregro Company (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.
2. Get to a safe and reliable factor beyond the manufacturer's control prevent it from incurring all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those set forth by the manufacturer under 1 above) resulting from handling, storage and use of this material.

1202-170

U.S.D.A. Reg. No.