

## ETHION 4 DUST

#### DIRECTIONS FOR USE

APPLES: Apply 50-90 lbs. per acre to control European red mite, two-spotted mite, brown almond mite, apple rust mite. Do not apply more than twice during the fruiting period or within 40 days of harvest. CAUTION--do not use on wealthy or melba varieties.

GRAPES: Apply 15-25 lbs. per acre to control grape leafhoppers two-spotted mites. Willamette mites. Do not apply within 15 days of harvest if not dusted more than twice during fruiting period or do not apply within 30 days if applied not more than 3 times during fruiting season.

PEACHES: Apply 40-60 lbs. per acre to control European red mite, two-spotted mite, brown almond mite. Do not apply within 30 days of harvest. Do not apply more than twice during fruiting season.

BEANS: (Succulent and Dry): Apply 20 lbs. per acre to control bean leaf beetle, Mexican bean beetle, two-spotted mite. Do not apply within 2 days of harvest. Do not feed treated bean foliage to livestock.

COTTON: Apply 25-40 lbs. per acre to control spider mite, lygus bug. Do not apply after bolls open. Do not graze dairy or meat animals in treated fields.

MELONS: Apply 30-40 lbs. per acre to control melon leafhopper, melon leafminer, serpentine leafminer, two-spotted mite. Do not apply within 2 days of harvest.

STRAWBERRIES: Apply 20-25 lbs. per acre to control two-spotted mite, strawberry leaf roller. Do not apply within 2 days of harvest.

Lot No.

4/70CN

ACTIVE INGREDIENTS:

\*Ethion.....4.00%

INERT INGREDIENTS:.....96.00%

100.00%

\*0, 0, 0', 0'-tetraethyl S, S'-methylene bisphosphorodithioate

U. S. Patent Number: 2,873,228

# CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May be absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs.



NET WEIGHT

LBS.

ACCEPTED

SEP 2 4 1970

UNDER THE FEDERAL INSECTICIDE FUNCICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON RECEIVED.

ED UNDER NO. 1202-185

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Apply this product only as specified on this label. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

CONDITIONS OF SALE: 1. PureGro Company (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond the manufacturer's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under I above) resulting from handling, storage, and use of this material.

USDA NO. 1202-183



### ETHION 4 DUST



#### DIRECTIONS FOR USE

APILES Apply 5 % lts, per acre to control Foregean ted mite, two spotted mite, brown almost note, apple rust made. Do not apply more than twice forms the fruiting period or within 40 may of harvest. PARTION--do not use on wealthy in

GRAPES: Apply 15-25 lbs. per sore to mitril grape leaft, ipers two-spatted mitrs willaments mites. Do not apply within 15 lays of harvest if it tusted over than twice during fruiting term is or do not apply within 30 days it applied to mire than 3 times during fruiting season.

PEACERS Apply 40-rulbs, per abre to dorth. European red mite, two-spotted mite, brown aumoud mate. Do not apply within 30 days of harvest. Do not apply more than twice further front in measurements.

COTTON Apply 25-4 list one are to the bound of the country of the control of the country of meat are made in treated fields.

MRID Apply 30-4 to per air of trico leafhopper, melon leafmaner semperator leafmaner two-spected mater. To our airle within 1800 to narvest.

Let No.

A TOTAL INCHES I PATS:

T. Fatert Number, 2,873,228

## CAUTION KEEP OUT OF REACH OF CHILDREN

High for an aware seed. May be absorbed the uight with a contract of a contract with a contrac



NET WELLET 136.

ACCEPTED

SEP 2 4 1970

THE STATE AND THE PARTICIDE ACT FOR ICON MICE AND THE PARTICIDE ACT FOR ICON MICE AND THE PARTICIDE ACT FOR ICON MICE AC

This product is foxic to fish and wildlife. Keep out of lakes, streams or jonds. Apply this product only as specified on this label. To not apply when wither inditions favor drift from areas treated. The not apply where rimoff is likely to occur. Do not contaminate water by leaning of equipment, or disposal of wastes.

This product is toxic to fees and should not be applied when tees are actively visiting the area.

Do not reuse emity container. Lestroy it by furgion with waste or burbles. This may from on kellin tumes.

that the material control of the form of the facturer is organish that the material control of the facturer is organish to the facturer of the facturer of the facturer materials that is a control of the facturer of the fac

The first of the property of t

USDA N. 1.02-183