

## SEVIN® BAIT PELLETED

ACTIVE INGREDIENT

Carbary1 (1-naphthyl. N-methylcarbamate)......5%

INERT INGREDIENTS

U.S. Pat. Nos. 2,903,478 --3,009,855

Trade Mark of Union Carbide Corporation for the . active ingredient 1-naphthyl N-methylcarbamate.

## KEEP OUT OF REACH OF CHILDREN

HAPMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid Breathing Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid prolonged and repeated contact with skin. Wash Hands and Face Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily.

Note for Physicians: Carbaryl is a moderate reversible, Cholinesterase inhibitor. Atropine is antidotal. AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

LOT NO. 4/70CN

> USDA Reg. No. 1202-185

NEM WETCHM



ACCEPTED

UNDER THE FEDERAL INSECTICION FUNGICIDE AND RODENTICIDE ACT ECONOMIC POMON REGISTRE,

DEC 1 6 1970

## DIRECTIONS FOR USE

Broadcast applications may be made with either ground equipment or aircraft. Repeat treatment as necessary.

For Cutworms, Armyworms, Sowbugs, Grasshoppers, Darkling Ground Beetles and Crickets

Cucumbers, Squash, Melons -- 20 lbs. per acre. May be applied up to harvest.

Alfalfa, Cotton, Peas -- 30 lbs. per acre. May be applied up to harvest on Alfalfa and Peas. Do not apply within 7 days of harvest or grazing on cotton.

Asparagus, Beans, Broccoli, Beets, Brussels Sprouts, Cabbage, Carrots, Cauliflower, Collards, Corn, Eggplants, Endive, Lettuce, Horseradish, Kale, Okra, Parsley, Parsnips, Spinach, Strawberries, Sugar Beets, Swiss Chard, Tomatoes, Turnips -- 40 lbs. per acre. Apply during cutting season on Asparagus. Do not repeat application on Asparagus within 3 days and do not apply within 1 day of harvest. Do not apply within 1 day of harvest on Strawberries. Do not apply within 3 days of harvest on Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Head Lettuce, Horseradish and Parsnips. Do not apply within 14 days of harvest on Sugar Beets, Collards, Endive, Kale, Leaf Lettuce, Parsley, Spinach and Swiss Chard. Do not apply to Beets or Turnips within 3 days of harvest or within 14 days if tops are to be used for food or feed.

Grapes -- 25 to 50 lbs. per acre. May be applied up to harvest.

CONDITIONS OF SALE: 1. PureGro Company (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond the manufacturer's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, A damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer