

DIRECTIONS FOR USE

To apply, mix required amount of THIODAN 2 C O with sufficient water to cover 1 acre. (By ground apply a minimum of 20 gallons per acre on vegetable or row crops. By air, apply a minimum of 3 to 10 gallons per acre). Apply with continuous agitation. Do not plant any root crops other than carrots, potatoes, and sweet potatoes as follow up crops.

Apply when insects first appear and repeat as required to maintain effective control. Coverage of both upper and lower leaf surfaces is essential for good control. Observe use limitations. If insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides. Observe days interval between last application and harvest indicated by number in parenthesis following crop.

SUCCULENT BEANS, EXCEPT LIMAS: Mexican Bean Beetle -- 1 qt. per acre. Do not apply after pods begin to form. Do not feed treated threshings to livestock. Do not apply more than 3 times per season. Do not allow livestock to graze in treated fields.

BROCCOLI (7), CABBAGE (7): Cabbage Aphid, Cabbage Looper, Imported Cabbage Worm, Diamondback Moth Larvae, Cross-striped Cabbage-worm, Flea Beetles, Harlequin Bugs, Stink Bugs -- 3 to 4 pts. per acre.

CAULIFLOWER: Cabbage Looper, Imported Cabbage Worm, Diamondback Moth Larvae, Cross-striped Cabbage worm, Flea Beetles, Harlequin Bugs, Stink Bugs -- 1 1/2 qts. per acre. Do not apply after edible parts begin to form. Close trim outer leaves prior to marketing.

CUCUMBERS (0), SQUASH (SUMMER & WINTER) (0): Aphids, Cucumber Beetle, Pickleworm, Melon worm, Squash Beetle, Squash Bug, Squash Vine Borer, Striped Cabbage Flea Beetles -- 1 to 2 qts. per acre. For Vine Borer control, apply weekly to flower buds, stems and vines beginning when moths first appear.

PEPPERS (1): Flea Beetle, Green Peach Aphid, Pepper Maggot, Hornworms -- 1 qt. per acre.

POTATOES (0): Aphids, Colorado Potato Beetle, Plant Bugs (including Shield Shaped Bugs), Potato Flea Beetle, Potato Leafhopper, Potato Tuberworm, Southern Armyworm, Whitefly, Green Stinkbug -- 1 to 2 qts. per acre. European Corn Borer, Potato Psyllid -- 1 1/2 to 2 qts. per acre. False Chinch Bug -- 2 qts. per acre. Three lined Potato Beetle -- 1 qt. per acre.

EGGPLANTS (1): Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid, Green Stink Bug, Whitefly -- 1 qt. per acre.

LETTUCE, HEAD (14): Cabbage Looper, Green Peach Aphid, Diamondback Moth Larvae, Imported Cabbage Worm -- 3 to 4 pts. per acre. Do not make more than 3 applications after thinning. Remove wrapper leaves at harvest. Do not feed crop refuse to livestock.

MELONS (0): Aphids -- Use 1 qt. per acre. Cucumber Beetles, Striped Cabbage Flea Beetle, Melonworm, Pickleworm, Squash Beetle, Squash Bug, Squash Vine Borer -- 1 to 2 qts. per acre. For Vine Borer control, apply weekly to flower buds, stems, and vines beginning when moths first appear. Cabbage Looper, Omnivorous Leaf Roller -- 2 qts. per acre.



THIODAN 2 C O INSECTICIDE WITH VEGETABLE OIL

ACTIVE INGREDIENTS:

Endosulfan (Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide)	23.7%
Cotton Seed Oil	46.0%
Petroleum Distillates	24.9%

INERT INGREDIENTS

5.4%
100.0%

E.P.A. REG. NO. 1202-188-AA

WARNING

KEEP OUT OF REACH OF CHILDREN

(See Additional Warning on Right Panel)

DO NOT USE IN UNDILUTED FORM

* Thiodan is a registered trademark of Canadian Hoechst Limited.



LOS ANGELES, CALIFORNIA

NET CONTENT

GALS

TOMA
rado
fly --
Fruits
2 qts
Loope

COND
warran
tion o
on. M
repres
MERC
imple

2. Cri
contro
the us
to, da
plied,
to oth
product
thought
knowle
assum
handli

lot No

NOT FO



THIODAN 2 C O
INSECTICIDE WITH VEGETABLE OIL

ACTIVE INGREDIENTS:

Endosulfan (Hexachlorocyclohexa- hydromethano 2,4,3 benzodioxathiepin oxide)	23.7%
Cotton Seed Oil	46.0%
Petroleum Distillates	24.9%

INERT INGREDIENTS

5.4%
100.0%

E.P.A. REG. NO. 1202-188 AA

WARNING

KEEP OUT OF REACH OF CHILDREN

(See Additional Warning on Right Panel)

DO NOT USE IN UNDILUTED FORM

* Thiodan is a registered trademark of Canadian Hoechst Limited.



LOS ANGELES, CALIFORNIA

NET CONTENT

GALS

TOMATOES (Field and Greenhouse) (1): Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid, White fly - 1 qt. per acre. Hornworms - 1 to 2 qts. per acre. Tomato Fruitworm, Tomato Russet Mite, Yellow striped Armyworm - 2 qts. per acre. Stink Bugs - 1 1/2 to 2 qts. per acre. Cabbage Looper - 1 1/2 qts. per acre

CONDITIONS OF SALE: 1. PureGro Company (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit to use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond the manufacturer's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above) resulting from handling, storage, and use of this material.

Lot No. _____ 8 71 CN

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

WARNING: Hazardous if swallowed, inhaled, or absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking; wear clean clothing. During commercial or prolonged exposure in spray-mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type passed by the U.S. Bureau of Mines for Endosulfan protection. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

CAUTION: Workers entering treated fields within 24 hours should wear protective clothing.

WARNING! FLAMMABLE

Keep away from heat and open flame.
Keep container closed.

Leaking packages should be removed to a safe place.
Do not store at temperatures below 20° F.

This product is toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Do not apply to any area not specified on the label. Birds feeding on treated areas may be killed. This product is toxic to bees and should not be applied when bees are actively visiting the area.

Dealers Should Sell in Original Packages Only.

USAGE CAUTION: DO NOT ALLOW THIS MATERIAL TO DRIFT ON TO NEIGHBORING CROP OR NON CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH DIRECTIONS, BECAUSE PLANT INJURY, EXCESSIVE RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

