



DIAZINON[®] SULFUR 2-50 DUST

ACTIVE INGREDIENTS:

1,2-diethyl 0-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate	100.0
Sulfur	50.0
INERT INGREDIENTS	48.0
	100.0

E.P.A. Reg. No. 1202-189-AA

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May be absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid storage near food and feed products.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes and containers. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources.

This product is highly toxic to bees. Do not use for treatment of flowering crops. Protective information has been obtained from your representative. For full information, consult your representative.

Do not use for treatment of flowering crops. Do not use for treatment of flowering crops.

Do not use for treatment of flowering crops. Do not use for treatment of flowering crops.



APPLES: Apply 45-75 lbs. per acre to control apple aphids, apple leafhoppers, apple scale crawlers. Do not apply within 10 days of harvest.

CHERRIES: Apply 45-75 lbs. per acre to control cherry aphids, cherry fruit flies, cherry scale crawlers. Do not apply within 10 days of harvest.

GRAPES: Apply 40-50 lbs. per acre to control grape aphids, grape leafhoppers, grape berry moth, grape leaf folder. Repeat as necessary. Do not apply within 10 days of harvest.

PEACHES: Apply 45-75 lbs. per acre to control peach aphid, brown mite, two-spotted mite, San Jose scale crawlers, olive scale crawlers. Do not apply within 10 days of harvest.

POTATOES: Apply 30-100 lbs. per acre to control aphids, flea beetles, Colorado potato beetles, dipterous leaf miners, banded cucumber beetles, leafhoppers, Southern armyworms. Do not apply within 35 days of harvest. Foliage application.

TOMATOES: Apply 60-80 lbs. per acre to control aphids, dipterous leaf miners, banded cucumber beetles, fall armyworms, southern armyworms, vinegar fly. Do not apply within 3 days of harvest.

ALFALFA, CLOVER: Apply 24-30 lbs. per acre to control spotted alfalfa aphids, yellow clover aphids, pea aphids, leafhoppers, plant bug nymphs, etc. Do not apply within 7 days of cutting or grazing by and all being finished for slaughter.

BERRIES: Apply 60-80 lbs. per acre to control aphids, leafhoppers, etc. Do not apply within 3 days of harvest.

PEARS: Apply 45-75 lbs. per acre to control pear aphid, pear leafhopper, etc. Do not apply within 10 days of harvest.

PLUMS: Apply 45-75 lbs. per acre to control plum aphid, etc. Do not apply within 10 days of harvest.

RAISINS: Apply 45-75 lbs. per acre to control grape aphids, etc. Do not apply within 10 days of harvest.

STRAWBERRIES: Apply 45-75 lbs. per acre to control strawberry aphid, etc. Do not apply within 10 days of harvest.

WATERMELONS: Apply 45-75 lbs. per acre to control watermelon aphid, etc. Do not apply within 10 days of harvest.

ZUCCHINI: Apply 45-75 lbs. per acre to control zucchini aphid, etc. Do not apply within 10 days of harvest.

Oct 20, 1964
1202-189