

DIRECTIONS FOR USE

For application by ground or air equipment.

BEANS: Mexican Bean Beetle, Leafhoppers, Powdery Mildew--10 lbs. per acre. **LYGUS, STINK BUGS**--20 lbs. per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary repeat every 7 to 10 days. May be used up to harvest.

CORN: Corn Earworm--15 to 20 lbs. per acre. Apply at 3 day intervals starting when silks first appear. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations. Make 4 to 5 applications until silks dry or turn brown. May be applied up to harvest.

COTTON: Early-Season--Thrips, Cotton Fleahopper--8 to 15 lbs. per acre as soon as insects appear. Mid and Late-Season--Boll Weevil, Bollworm, Pink Bollworm, Cotton Leafworm, Lygus Bugs, Cotton Leaf Perforator--20 lbs. per acre. Apply when insects appear and repeat at 5 to 7 day intervals. Aphid populations will usually be suppressed when this insecticide is used on a regular schedule of treatments.

GRAPES: Grape Leaf Folder--1st Brood: 10 to 15 lbs. per acre just before first brood larvae emerge from the first rolls. 2nd and 3rd Brood: 10 to 20 lbs. per acre. Grape Leafhopper, Powdery Mildew--10 lbs. per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary repeat every 7 to 10 days. Thorough coverage is essential. May be applied up to harvest. This application may be expected to control DDT-resistant leafhoppers.

PEACHES, NECTARINES: Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Cat-Facing Insects, Periodical Cicada, Powdery Mildew, Brown Rot--35 to 40 lbs. per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary repeat every 7 to 10 days. Do not apply 1 day before harvest for peaches and 3 days before harvest for nectarines.

PLUMS, PRUNES, CHERRIES: Fruit Tree Leaf Roller, Peach Twig Borer, Plum Curculio, Red-banded Leaf Roller, Codling Moth, Eastern Tent Caterpillar, Orange Tortrix, Western Tussock Moth--40 to 50 lbs. per acre. Apply when insects appear and repeat as necessary. Make full coverage application. Do not apply within 1 day of harvest.

POTATOES, EGGPLANT, PEPPER: Colorado Potato Beetle--10 lbs. per acre. Leafhopper, Flea Beetle, Lygus Bugs--20 lbs. per acre. Apply when insects appear. Apply dust thoroughly to upper and lower leaf surfaces. If needed, repeat applications at 7 to 10 day intervals.

TOMATOES: Tomato Fruitworm, Fall Armyworm, Flea Beetles, Tomato Hornworm, European Corn Borer, Stink Bugs--30 to 40 lbs. per acre. Apply thoroughly at first signs of insect infestation. Repeat as necessary. May be applied up to the day of harvest. Because of the sulfur content of this dust, first obtain permission from your canner or processor before using it on tomatoes intended for sale to such canner or processor.

LEAFY VEGETABLES AND ROOT CROPS: (Carrots, Chinese Cabbage, Collards, Dandelion, Endive, Beet Horseradish, Kale, Lettuce, Mustard Greens, Parsley, Parsnip, Radish, Rutabaga, Salsify, Spinach, Swiss chard, Turnip): Apply 10-20 lbs. per acre to control flea beetles, leafhoppers, harlequin bug, six-spotted leafhopper (aster yellows vector), corn earworm, armyworms, lygus and stink bugs. Do not apply 14 days before harvest.

SEVIN® SULFUR DUST 10-50

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)	10.0%
Sulfur	50.0%
INERT INGREDIENTS:	40.0%
	100.0%

Fenage Sulfur

CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing dust. Avoid contact with skin; wash thoroughly after using. Avoid storage near feed and food products.

Do not apply to cucurbits unless the variety is known to be sulfur tolerant under local conditions. Avoid drift to sulfur sensitive plants.

Do not apply to any area not specified on the label.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

Registered trademark of Union Carbide Corporation for the active ingredient (Carbaryl).

BURN EMPTY CONTAINER
KEEP OUT OF SMOKE



Lot No. _____ 1/71CN

Net Contents _____ lbs.

ACCEPTED

MAR 22 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 1202-190

ASPARGUS: Apply 15-20 lbs. per acre to control asparagus beetle. Apply during cutting season. Do not repeat application within 3 days. Do not apply 1 day before harvest.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI: Flea beetle, harlequin bug, corn earworm, armyworms. For small loopers after heads form, use SEVIN every 5 to 7 days--Apply 15-20 lbs. per acre. Do not apply 3 days before harvest.

FILBERT: Filbert worm, filbert aphid, filbert leaf roller--Apply 40 lbs. per acre. No time limitation.

FORAGE AND FEED CROPS

ALFALFA, CLOVERS, PEANUTS, SOYBEANS, COMPEAS: Blister beetles, Mexican bean beetle, thrips, leafhoppers, three-cornered alfalfa hopper, cucumber beetles, bean leaf beetle, Japanese beetle, alfalfa caterpillar, velvetbean caterpillar and green cloverworm, armyworms, webworms, corn earworm, grasshoppers. Apply 10 lbs. per acre. No time limitation.

SORGHUM: (Grain and Forage): Armyworm, webworm, corn earworm, stink bugs, sorghum midge, (direct dust into forming heads. Apply three to four days after heads have emerged from boot.), grasshoppers--Apply 20 lbs. per acre. Do not apply 21 days before harvest of grain. No time limitation on sorghum forage.

RICE, FORAGE GRASSES, PASTURE: Armyworms, thrips, apply 15 lbs. per acre. For optimum thrips control in grasses grown for seed, direct dust into boot. Do not apply to Rice within 14 days of harvest. May be applied up to harvest on Pastures and Forage Grasses.

CONDITIONS OF SALE: 1. PureGro Company (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond the manufacturer's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above) resulting from handling, storage, and use of this material.

U.S.D.A. Reg. No. 01202-00190-AA-00000