

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals
May cause irritation of nose, throat, eyes and skin. Do not breathe dust or vapor. Use **PROTECTIVE CLOTHING** consisting of long pants, long sleeves shirt, gloves, hat and boots made to wear during mowing and loading.

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 2 glasses of water to drink and call a physician. Never give anything by mouth to an unconscious person.

IF INHALED: Move subject to fresh air.

IF IN EYES: Flush eyes with large amounts of water for at least 15 minutes. Consult a physician if irritation persists.

IF ON SKIN: Wash affected skin areas with soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not enter treated areas without protective clothing until dusts have settled.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or Oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: "WARNING: Area to be treated with Dithane on (date of application). Do not enter without appropriate protective clothing. All dusts have settled. If accidentally exposed, consult a physician, or Poison Control Center."

STORAGE & DISPOSAL

Do not contaminate water, food, feed by storage and disposal.

STORAGE: Store in a cool, dry place. Keep away from fire and sparks. Do not allow to become wet or over-heated in storage. This may bring on chemical changes which will impair the fungicidal effectiveness and may also generate flammable vapors. Keep container closed when not in use.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at any approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

LOT NO. _____ 8/84 C.D.

M-45 DUST
CONTAINS *DITHANE M-45
ACTIVE INGREDIENT:

Zinc ion and manganese ethylene bisdithiocarbamate	3.2%
a coordination product of manganese	0.64%
Zinc	0.08%
ethylene bisdithiocarbamate	2.48%
INERT INGREDIENTS96.8%
		100.0%

E.P.A. Reg. No. 1202-191-AA

E.P.A. Est. No.

U.S. Pat. Nos. 2,504,404; 2,974,156 and Patent Pending.

Registered Trademark of Rohm & Haas.

CAUTION KEEP OUT OF REACH OF CHILDREN

ACCEPTED

FEB 4 1985

Under the Federal Insecticide,
Fungicide and R伯onticide Act,
as amended, for the pesticide
registered under
EPA Reg. No 1202-191

Distributed

By



KENT SACRAMENTO, CA 95814

NET CONTENTS _____ LBS.

CROP RECOMMENDATION

—rates given in pounds per acre. To be applied by ground duster or aircraft. Application should begin early in the season. Treatments should be repeated whenever as needed, generally every 7-10 days and continued as permitted by the limitations listed for each crop.

FRUIT AND NUT CROPS:

APPLES, PEARS, CRABAPPLES, QUINCE: 240 lbs scab, cedar-apple rust, black rot, fly speck, sooty blotch, bitter rot, brown rot. Do not apply within 15 days of harvest. Do not graze livestock in treated areas.

GRAPES: (California Only) 50-100 lbs bunch rot, deadarm, early season downy mildew. Do not apply after fruit set.

CRANBERRIES: 60-100 lbs fruit rot. Do not apply within 30 days of harvest.

PAPAYAS: 50-80 lbs anthracnose, phytophthora fruit rot. No time limitation.

FIELD CROPS:

BARLEY, OATS, RYE, WHEAT: 50 lbs. helminthosporium leaf blight, septoria leaf spot. Do not apply within 26 days of harvest. Do not graze livestock in treated area within 26 days after application. Do not apply more than 3 applications in a season.

COTTON: 45-50 lbs rust (Southwest only). Do not apply after bolls are open. Do not graze livestock in treated areas or feed gin trash to livestock.

PEANUTS: 30-35 lbs cercospora leaf spot. Do not apply within 14 days of harvest. Do not feed vine hay to livestock.

VEGETABLE CROPS:

ASPARAGUS: 50 lbs rust. Apply after harvest to ferns only.

CARROTS: 30-50 lbs. cercospora and alternaria leaf spot. Do not apply within 7 days of harvest. Do not use treated tops for food or feed purposes.

CELERY: 50 lbs. early blight, late blight. Do not apply within 7 days of harvest.

CUCUMBERS, CANTALOUPE, HONEYDEWS, MUSK-MELONS, SUMMER SQUASH, WATERMELONS: 50-75 lbs downy mildew, anthracnose, scab, alternaria leaf spot, cercospora leaf spot, gummy stem blight. No time limitation.

POTATOES: 25-50 lbs late blight, early blight. No time limitation.

TOMATOES: 35-75 lbs late blight, early blight, gray leaf mold, anthracnose. No time limitation.

PUREGRO CONDITIONS OF SALE AND WARRANTY

PureGro Company warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label. PureGro makes no other warranty, express or implied, of fitness or merchantability nor authorizes any agent or representative to make any variation or exception from this warranty.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of PureGro Company. Buyer assumes all risk of use, storage, or handling of this material not in strict accordance with directions given herewith.