

ACCE

Aug 18-1971

ORDER 1207-197



TERRACHLOR 20 DUST

DIRECTIONS FOR USE

BEANS: Band treatment--Apply 5-20 lbs. per 8,400 linear feet of row for pole beans or 14,500 linear feet for bush beans to control root and stem rots (rhizoctonia). Apply a 6 to 12 inch wide band over seed and surrounding soil at planting time. Do not feed treated vines to livestock.

CANES: Apply 20-25 lbs. per 8,400 or 14,500 linear feet of row depending on growth type. Apply to base of canes when disease first appears but not later than first harvest. Do not cultivate after application. May be applied as a pre-harvest application to plants and soil surface when disease first appears but not after first bloom. Do not feed after application. Do not feed treated vines to livestock.

BEETROOT, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER: Apply 20-30 lbs. per 13,100 linear feet of row in 12-15 inch bands, rototilled to a depth of 4 to 6 inches in a preplant operation; or 150-300 lbs. per acre in a preplanting broadcast application rototilled or disked and cross-disked to a depth of 4 to 6 inches to control clubroot (plasmodiophora), wire stem or black rot (corticium solani) 25 lbs. per 13,100 linear feet of row. Apply in an 8 inch band centered on row prior to seeding. Mix into top 2 inches of soil.

COTTON: Apply 10 lbs. per 12,400 linear feet of row to control rhizoctonia induced damping-off, seedling blight, seedling rot, sore-shin. Apply in furrow over seed and surrounding soil at planting time or 3-5 lbs. per acre mixed with seed in planter-box. Do not allow feeding or grazing of cotton foliage by livestock. Do not plant any root crop not registered for PCNB in rotation on PCNB treated soil.

PEPPERS: Apply 35 to 37-1/2 lbs. per 14,500 linear feet of row to control Southern blight. Dust open "V" trench prior to setting transplants.

ACTIVE INGREDIENTS:
Pentachloronitrobenzene 20
INERT INGREDIENTS: 80
100%

EPA Reg. No. 1202-00192-AA

CAUTION KEEP OUT OF REACH OF CHILDREN

Avoid breathing of dust. This material may cause skin irritation. Wash hands and face thoroughly after using. Wear protective clothing and a respirator when loading the spreader or duster. Avoid contamination of feed and foodstuffs. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.



NET WEIGHT _____ LBS.

TOMATOES: (Field grown): Apply 35-37 1/2 lbs. per 7,300 linear feet of row for nonstaked plants or 10,900 linear feet for staked plants to control Southern blight. Dust opened "V" trench prior to setting transplants.

ROSES: Botrytis Storage Rot-Stack dormant roses "root to root" and apply by liberally dusting branches immediately after stacking.

CONDITIONS OF SALE: 1. PureGro Company manufacturer warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond the manufacturer's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above) resulting from handling, storage, and use of this material.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

Lot No. _____ 771CN