



GAVIDOX SULFUR 5-50

DIRECTIONS FOR USE

Apply the specified amount of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infestation and repeat at regular intervals when needed to maintain control, but observe use limits given on label.

RECOMMENDATIONS

Field Crops

ALFALFA: Apply 20 lbs. per acre to control alfalfa caterpillar, alfalfa webworm, Western yellow-striped armyworm, beet armyworm, variegated cutworm, lygus bugs, stink bugs, tarnished plant bug. 1 application per cutting. Do not apply within 14 days of cutting or grazing.

COTTON: Apply 20-30 lbs. per acre to control cotton fleahopper, cotton leafworm, Western yellow-striped armyworm, beet armyworm, Southern garden leafhopper, black fleahopper complex, cotton leaf perforator, leaf roller, lygus bugs, stink bugs. Do not pasture treated fields within 14 days of application. Do not apply within 7 days of picking. Dylox applications may occasionally cause marginal leaf burn.

SAFFLOWER: Apply 25-30 lbs. per acre to control armyworms, dipterous leaf miners, imported cabbageworm, lygus bugs, stink bugs, thrips, variegated cutworm. Do not apply after flowering stage. Do not apply more than twice per year.

SUGAR BEETS: Apply 20-30 lbs. per acre to control beet webworm, variegated cutworm, dipterous leaf miners, alfalfa webworm, beet armyworm. Apply 30 lbs. per acre to control salt marsh caterpillar. Do not apply within 14 days of harvest. Tops harvested less than 28 days after treatment do not feed to livestock.

Seed Crops Only

CLOVER, SOYBEANS: Apply 25 lbs. per acre to control armyworms, dipterous leaf miners, imported cabbageworm, lygus bugs, stink bugs, variegated cutworm. Do not pasture or use treated crop for feed, food or forage.

Vegetables

BEANS (Dry and Snap): Apply 25-30 lbs. per acre to control Western bean cutworm, armyworms, dipterous leaf miners, imported cabbage worm, lygus bugs, Mexican bean beetle, stink bugs, variegated cutworm. Do not apply within 14 days of harvest. Do not use treated vines as food for dairy or meat animals.

IMA BEANS: Apply 20-30 lbs. per acre to control Western bean cutworm, armyworms, dipterous leaf miners, imported cabbage worm, lygus bugs, Mexican bean beetle, stink bugs, variegated cutworms. Do not apply more than twice per season. Do not harvest for food or forage within 30 days of treatment.

COMPEAS (Southern Peas, Blackeyed Peas, Crowder Peas): Apply 25-30 lbs. per acre to control armyworms, dipterous leaf miners, imported cabbageworm, lygus bugs, variegated cutworm. Apply only twice per season. Do not apply within 14 days of harvest. Do not use vines as food for dairy or meat animals.

BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER: Apply 20 lbs. per acre to control Western yellow-striped armyworm, imported cabbageworm, diamondback moth. Do not apply within 21 days of harvest.

ACTIVE INGREDIENTS:

Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate*..... 5%
Sulfur..... 50%

INERT INGREDIENTS.....

45%
100%

CONTAINS 5% DYLOX

Dylox is a reg. U.S. Pat. Office by Farbenfabriken Bayer, A. G. Chemagro Corporation Licensee

U.S. Patent No. 2,701,224

WARNING!

KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes or on skin. Do not breathe dust. Always wash thoroughly with soap and warm water after handling. Wash contaminate clothing with soap and hot water before re-use.

TO PHYSICIAN: Atropine sulfate is antidotal 2-PAM is also antidotal and may be administered in conjunction with atropine.

E.P.A. Reg. No. 01202-194-AA

RESTRICTIONS


Do not use on other crops used for food or forage. Use only according to label directions. Application at rates above those shown may result in illegal crop residues. Do not treat food crops grown in greenhouses.

Do not contaminate food or feed.

If poisoning occurs, obtain prompt medical aid.

Do not reuse container. Destroy when empty.

Lot No. _____ 8/71CN



NET WEIGHT _____ LBS.

CARROTS: Apply 20-30 lbs. per acre to control Western yellow-striped armyworms, imported cabbageworm, variegated cutworm, diamondback moth, dipterous leaf miners, beet armyworm, lygus bugs. Repeat as necessary. Do not use tops for food or feed. Do not apply within 28 days of harvest.

COLLARD, LETTUCE: Apply 20 lbs. per acre to control variegated cutworm, armyworms, beet webworm, diamondback moth, dipterous leaf miners, salt-marsh caterpillar. Repeat as necessary until heads begin to form. Do not apply within 28 days of harvest.

PEPPERS: Apply 20-25 lbs. per acre to control dipterous leaf miners, pepper maggot, serpentine leaf miners. Repeat as necessary. Do not apply within 21 days of harvest.

PUMPKIN: Apply 25 lbs. per acre to control variegated cutworm, squash bug. Apply only once per season and not within 14 days of harvest.

TABLE BEETS: Apply 20-30 lbs. per acre to control variegated cutworm, alfalfa webworm, beet webworm, dipterous leaf miners, salt-marsh caterpillar, beet armyworm, lygus bugs. Repeat as necessary. Do not apply within 28 days of harvest. Do not eat tops.

TOMATOES: Apply 20-25 lbs. per acre to control serpentine leaf miners, tomato hornworms, dipterous leaf miners. Repeat as necessary. Do not apply within 21 days of harvest.

ORNAMENTALS

FLOWERS, SHRUBS, TREES: Apply 20-30 lbs. per acre to control armyworms, dipterous cutworms, dipterous leaf miners, lygus bugs, stink bugs, tarnished plant bug, tobacco budworm, webworms. CAUTION: Phytotoxicity has occurred on certain varieties of carnations and zinnias. When large plantings of these flowers are to be dusted for the first time, treat only a few plants and observe for 4 to 5 days before dusting the entire planting.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water.

Do not apply when weather conditions favor drift from areas treated.

Do not apply where runoff is likely to occur.

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

CONDITIONS OF SALE: 1. PureGro Company (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond the manufacturer's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though the label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above) resulting from handling, storage, and use of this material.