

DIRECTIONS FOR USE

VEGETABLE CROPS:

ASPARAGUS: Apply 15-25 lbs. per acre to control asparagus beetle, rust. Begin after cutting stops. Repeat at 7 to 10 day intervals through season.

BEANS: Apply 15-25 lbs. per acre to control Mexican bean beetle, leaf hoppers, bean leaf beetle, cucumber beetle, western bean cutworm, flea beetle, Japanese beetle, velvetworm, leaf spot, powdery mildew, rust. Begin when first true leaves form, or at first sign of disease. Repeat at 7-10 day intervals.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KALE, TURNIPS, COLLARDS, KUTABAGA: Apply 15-25 lbs. per acre to control flea beetle, harlequin bug, corn earworm, armyworm, powdery mildew. Begin at first sign of disease. Repeat at 7-10 day intervals. Do not apply within 3 days of harvest. Do not treat turnips within 14 days of harvest if tops are to be used for food or feed. Do not apply within 14 days of harvest on collards and kale.

CARROTS: Apply 20-25 lbs. per acre to control flea beetles, leafhoppers, six-spotted leafhopper (aster yellows vector), armyworms, powdery mildew. Begin at first sign of disease. Repeat at 7 to 10 day intervals. Do not apply within 14 days of harvest.

EGGPLANT, PEPPERS, TOMATOES: Apply 24-53 lbs. per acre to control tomato fruitworm, fall armyworm, tomato hornworm, European corn borer, lace bug, Colorado potato beetle, flea beetles, leafhoppers, powdery mildew. Eggplants: Begin when first true leaves appear. Repeat at weekly intervals. Peppers: Begin when disease first appears. Repeat at weekly intervals. Tomatoes: For control of insects only.

LETTUCE: Apply 20-25 lbs. per acre to control flea beetles, leafhoppers, harlequin bugs, six-spotted leafhopper (aster yellows vector), corn earworm, armyworms, powdery mildew. Begin at first sign of disease. Repeat as necessary. Do not apply within 14 days of harvest.

PEAS: Apply 15-20 lbs. per acre to control leafhoppers, Colorado potato beetle, armyworm, grasshoppers, powdery mildew, rust. Begin early in season before disease appears. Repeat at 7 to 10 day intervals.



SEVIN SULFUR DUST

7.5-50

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)	7.5%
Sulfur	50.0%
INERT INGREDIENTS	42.5%
	100.0%

ACCEPTED
12/27/72
CAUTION
KEEP OUT OF REACH OF CHILDREN
1202-195

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin; wash thoroughly after using. Avoid storage near feed and food products.

This product highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

Sulfur may burn foliage when temperatures are excessively high.

Registered trademark of Union Carbide Corporation for the active ingredient (Carbaryl).

E.P.A. Reg. No. 1202-195-AA

Do not reuse empty container, destroy it by burning or burying with waste. Stay away from smoke or fumes.



NET WEIGHT LBS.

POTATOES: Apply 15-25 lbs. per acre to control Colorado potato beetle, flea beetles, leafhoppers, powdery mildew. Foliage application only. Apply before disease occurs, where powdery mildew occurs, repeat as necessary.

SUGAR BEETS: Apply 15-25 lbs. per acre to control leafhoppers, armyworms, webworms, grasshoppers, rust (puccinia). Apply at first sign of disease. Repeat at 7 to 10 day intervals. Do not apply within 14 days of harvest.

FIELD CROPS:

ALFALFA: Apply 12-21 lbs. per acre to control blister beetles, Mexican bean beetle, thrips, leafhoppers, three-cornered alfalfa hopper, cucumber beetles, bean leaf beetle, Japanese beetle, alfalfa caterpillar, velvetbean caterpillar, green cloverworm, armyworms, webworms, corn earworm, stink bugs, grasshoppers, cutworms, powdery mildew, rust leaf spot. Apply to stubble or when crop is not over 3 to 6 inches high or when diseases appear. Repeat at 7 to 14 day intervals. Limits 21 pounds per acre.

COTTON: Apply 25-33 lbs. per acre to control thrips, flea beetle, fleahopper, leafworm, striped blister beetle, boll weevil, bollworm, fall armyworm, leaf perforator, leaf rollers, leafhoppers, tarnished plant bug, light to moderate numbers of Western lygus bugs (aphids will be repressed by repeated applications in a schedule), pink bollworm, salt marsh caterpillar, stink bugs, grasshoppers. Begin when first true leaves appear. Limits 33 pounds per acre.

SOYBEANS: Apply 13-1/4 to 20 lbs. per acre to control blister beetles, Mexican corn beetle, thrips, leafhoppers, three-cornered alfalfa hopper, cucumber beetles, bean leaf beetle, Japanese beetle, alfalfa caterpillar, velvetbean caterpillar, green cloverworm, armyworms, webworms, corn earworms, stink bugs, grasshoppers, cutworms, powdery mildew, leaf spot. Begin when disease first appears. Repeat at 7-10 day intervals. Limits 20 pounds per acre.

CONDITIONS OF SALE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

LOT NO. 11/72CN