DIRECTIONS FOR USE

For aircraft or ground rig application. Apply recommended rates when disease appears and repeat at 7-10 day intervals as long as necessary.

FRUIT AND NUT CROPS

ALMONDS (California): Apply 30-130 lbs. per acre to control shot hole, brown rot. Do not apply later than 5 weeks after petal fall. Do not feed almond hulls to dairy animals or animals being finished for slaughter.

APRICOTS (California): Apply 40-150 lbs. per acre to control brown rot, shot hole. Do not apply within 14 days of harvest. NECTARINES, PEACHES (California): Apply 40-100 lbs. per acre to control shot nole, peach lear curl, fruit brown rot. Do not apply within 14 days of harvest. Remove residue by brushing.

VEGETABLE, FIELD AND FORAGE CROPS

BEANS (Dry): Apply 20-55 lbs. per acre to control downy mildew, rust. No time limitation. BEANS (Succulent): Use recommended rates and diseases controlled under beans (dry). Do not apply within 4 days of harvest. BROCCOLI, BRUSSEL SPROUTS, CABBAGE CAULIFLOWER, KOHLRABI: Apply 25-30 1bs. per acre to control downy mildew, alternaris leaf spot. Do not apply within 7 days of harvest of cabbage, broccoli. CELERY: Apply 25-50 lbs. per acre to control early and late blight. Remove excess residues by stripping, trimming, and washing. CUCUMBERS, SQUASH, MELONS, PUMPKINS: Apply 25-40 lbs. per acre to control dcwny mildew, anthracnose, gummy stem olight. 14Ch - 400

_OT



MANEB 5.6 DUST

ACTIVE INGREDIENT:

Maneb (Manganous ethylene
bisdithiocarbamate) 5.6%
INERT INGREDIENTS: . . . 94.4%

 $100.0^{cr}_{\ c}$

CAUTION KEEP OUT OF REACH OF CHILDREN

May be absorbed through skin and throat. Do not breathe dust.

BURN EMPTY BAG STAY OUT OF SMOKE



NO.	5/71 CN- (**)		
	hun 1)	NET WEIGHT	LBS

(Continued from left panel)

EGGPLANTS, PEPPERS: Apply 25-40 lbs. per acre to control fruit rot, anthracnose. LETTUCE: Apply 25-40 lbs. per acre to control downy mildew. Do not apply within 7 days of harvest. Remove residues from head lettuce by stripping and trimming, and from leaf lettuce by washing or other effective means. POTATOES: Apply 25-50 lbs, per acre to control early and late blight. SPINACH: Apply 25-45 lbs. per acre to control downy mildew, white rust. Do not apply within 7 days of harvest. Remove excess residue by washing. TOMATOES: Apply 25-70 lbs. per acre to control early and late blight, anthracnose, stemphylium, septora leaf spot, cladosporium grey mold. FLOWERS (Asters, Iris, Mums, Gladiolus): Apply 25-40 lbs. per acre to control stemphylium, curvularia, botrytis.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

CONDITIONS OF SALE: 1. PureGro Company (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material. 2. Critical and unforeseeable factors beyond the manufacturers control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above) resulting from handling, storage, and use of this material.

U.S.D.A. Reg. No. 01202-00200-AA-00000

CAUTION

KEEP OUT OF REACH OF CHILDREN

May be absorbed through skin and throat. Do not breathe dust.

BURN EMPTY BAG STAY OUT OF SMOKE



200

NSECTICIDE

HTICIDE ACE

WEIGHT LBS.

FLOWERS (Asters, Iris, Mums, Gladiolus): Apply 25-40 lbs. per acre to control stemphylium, curvularia, botrytis.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

CONDITIONS OF SALE: 1. PureGro Company (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material. 2. Critical and unforeseeable factors beyond the manufacturers control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above) resulting from handling, storage, and use of this material.

U.S.D.A. Reg. No. 01202-00200-AA-00000

CAUTION

KEEP OUT OF REACH OF CHILDREN

May be absorbed through skin and throat. Do not breathe dust.

BURN EMPTY BAG STAY OUT OF SMOKE



200

INSECTICIDE

NTICIDE ACT

WEIGHT LBS.

FLOWISS (Asters, Iris, Mums, Gladiolus): Apply 25-40 lbs. per acre to control stemphylium, curvularia, botrytis.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

CONDITIONS OF SALE: 1. PureGro Company (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material. 2. Critical and unforeseeable factors beyond the manufacturers control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above) resulting from handling, storage, and use of this material.

U.S.D.A. Reg. No. 01202-00200-AA-00000