

DIRECTIONS FOR USE

FOR AIRCRAFT: Add sufficient water to equal a minimum of 5 gallons of spray per acre.

GROUND SPRAYERS: Use recommended amount of PickQuik Liquid Defoliant in 20 to 60 gallons of water per acre for cotton.

COTTON: Apply 2 to 2½ gallons per acre by ground sprayer or aircraft. For pre-harvest defoliation and desiccation, six to eight nozzles per row are usually necessary to obtain good results with ground sprayer. Do not graze treated areas or feed gin waste to livestock. Do not apply within 7 days of harvest.

MILO, SORGHUM: 2-3 gallons per acre for harvest desiccation. Apply when fully mature, 7 to 10 days before harvest.

ENVIRONMENTAL CAUTIONS

Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Pick 2/12

**PICK - QUIK
LIQUID DEFOLIANT**

ACTIVE INGREDIENT:	
Sodium Chlorate	18.5%
INERT INGREDIENTS:	81.5%
	<hr/>
	100.0%

Contains Urea as a fire suppressant.

E.P.A. Reg. No. 1202-206-AA

**CAUTION
KEEP OUT OF
REACH OF CHILDREN**

Harmful if swallowed. Avoid skin contact. Wash clothing and spray equipment after handling. Avoid contamination of feed and foodstuffs. Keep opened containers away from children and animals. Keep children and animals off treated areas.

DO NOT USE THIS PRODUCT FOR ANY PURPOSE OTHER THAN THAT RECOMMENDED.

March 19, 1975
U.S. DEPARTMENT OF AGRICULTURE
FOR ECONOMIC POIS. AND PESTICIDES
ED UNDER NO. 211-100-100
NO ATTACHED COMMENTS.

NET CONTENT _____ GALS.

#1245

SUGGESTIONS FOR APPLICATION
IMMATURITY AND DROUGHT OF PLANTS ARE THE MOST FREQUENT CAUSES OF UNSATISFACTORY DEFOLIATION. Under most growing conditions proper maturity is indicated by the green foliage color becoming lighter and tinged with red, and by a definite reddening of the stems. Do not defoliate when plants are suffering from drought as evidenced by wilt condition of the plant. Cotton suffering from an alkali soil condition should not be defoliated. Thorough coverage of the plant is essential, whether applied by ground or air. Apply defoliant 10-21 days before picking date on cotton.

CONDITIONS OF SALE:

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

LOT NO. _____ 2/75 CN