

ACCEPTED

L141972

MIDLAND PESTICIDE
440 FLORENCE BLVD.
CORP. HOUSTON, TEXAS
77002

02-2417

DIRECTIONS FOR USE

Begin applications at first sign of infestation and repeat as necessary.
For use by either ground or aircraft.

Fruit and Nut Crops

WALNUTS: Apply 50-60 lbs. per acre to control aphids, spider mites. Do not apply within 10 days of harvest.

GRAPEFRUIT, LEMONS, ORANGES: Apply 40-50 lbs. per acre to control aphids, red spider mite, citrus cutworm, thrips. Do not apply within 7 days of harvest except in California and Arizona only 1 day.

PEACHES: (Western States Only) Apply 30-40 lbs. per acre to control aphids, fruit flies (Drosophila sp.), aphids, red spider mites, peach twig borer, lygus, thrips. Do not apply within 4 days of harvest.

STRAWBERRIES: Apply 40-50 lbs. per acre to control spider mites, aphids. Do not apply within 4 days of harvest.

Vegetables, Beans, Field Crops
(includes crops for seed, also)

BEANS
BEANS
BEAN

BLACK OY, Lima, green: Apply 4-5 lbs. per acre to control lygus, caterpillars, lygus bugs, thrips, leafhopper, and cutworms, loopers. Do not apply within 4 days of harvest.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, TOMATOES: Apply 4-5 lbs. per acre to control cabbage root fly, imported cabbage worm, diamond-back caterpillar, aphids. Do not apply within 1 day of harvest.

CARROTS, KANSAS: Apply 4-5 lbs. per acre to control cabbage root fly, imported cabbage worm, diamond-back caterpillar, aphids. Do not apply within 4 days of harvest.

CELERY: Apply 40-50 lbs. per acre to control loopers, aphids. Do not apply within 4 days of harvest.

CORN, LENTIL, SPROUT: Apply 40-50 lbs. per acre to control loopers, aphids. Do not apply within 1 day of harvest.

COTTON: Apply 35 lbs. per acre to control lygus, cotton leaf miner, armyworms, loopers, fleahoppers. Do not apply within 4 days of hand picking.

EGGPLANTS: Apply 40-50 lbs. per acre to control aphids, caterpillars. Do not apply within 1 day of harvest.

HOPS: Apply 25 lbs. per acre to control hop loopers. Do not apply within 4 days of harvest.

PEAS: Apply 40-50 lbs. per acre to control aphids, flea beetles. Do not apply within 4 days of harvest.

PEPPERS: Apply 20-25 lbs. per acre to control aphids, flea beetles. Do not apply within 1 day of harvest.

(continued on right panel)

Lot No. _____ 4/720N



**DIBROM SULFUR
4-40 DUST**

ACTIVE INGREDIENTS:	
Naled*	4
Sulfur	40
INERT INGREDIENTS	
	56
	100

*1,2-dibromo-2, 2-dichloroethyl dimethyl phosphate

U.S. Pat. No. 2,971,882

Registered by
T. M. of Chevron Chemical Company

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May cause irritation of eyes, nose and throat. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after using. Avoid contamination of feed and forklifts. Do not use on cherries as injury may result. Avoid drift to sulfur sensitive crops.

ADULT MOSQUITO CONTROL: Tidal Marshes, Swamps, Woodlands, Feed Lots and Pastures -- 8 to 10 lbs. per acre. Avoid farm buildings and dairy farms. Make applications during peak of infestation and repeat as necessary. Do not pasture animals or treated areas within 4 days after application. May be poisonous to fish and shellfish. Avoid use where they will be endangered.

(Continued on right panel)



NET WEIGHT _____ LBS.

(Continued from left panel)

TOMATOES: Apply 25 lbs. per acre to control two-spotted mites, aphids. Make applications during picking and continue at 5 to 10 days intervals when necessary. Do not apply to green tomatoes.

(Continued from front panel)

This product is toxic to fish and aquatic life. Areas may be killed. Keep out of areas likely to occur. Do not apply to water. Do not contaminate water. Apply this product only as directed.

Rinse equipment and containers thoroughly away from water supplies. Do not dump or bury with wastes.

This product is highly toxic to birds and mammals. Do not use in areas where birds and mammals are present.

Keep out of reach of children. Do not use on cherries as injury may result. Avoid drift to sulfur sensitive crops.

E.P.A. Reg. No. _____



DIBROM[®] SULFUR 4-40 DUST

ACTIVE INGREDIENTS:

Naled*	4%
Sulfur	40
INERT INGREDIENTS	56%
	100%

*1,2-dibromo-2, 2-dichloroethyl dimethyl phosphate

U.S. Pat. No. 2,971,882

R
T. M. of Chevron Chemical Company

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May cause irritation of eyes, nose, and throat. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after using. Avoid contamination of feed and rock luffs. Do not use on cherries as injury may result. Avoid drift to sulfur sensitive crops.

ADULT MOSQUITO CONTROL: Tidal Marshes, Swamps, Woodlands, Feed Lots and Pastures -- 8 to 10 lbs. per acre. Avoid farm buildings and dairy barns. Make applications during peak of infestation and repeat as necessary. Do not pasture animals on treated areas within 4 days after application. May be poisonous to fish and shellfish. Avoid use where they will be endangered.

(Continued on right panel)



NET WEIGHT _____ LBS.

(Continued from left panel)

TOMATOES: Apply 25 lbs. per acre to control fruit flies (*Drosophila* sp.), two-spotted mites, aphids. Make first application at 5 to 7 days before first picking and continue at 5 to 7 day intervals until harvest is complete or apply when necessary. Do not apply within 1 day of harvest.

(Continued from front panel)

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning (keep out of smoke).

This product should not be exposed to direct treatment or residues of a pesticide which may be obtained from your Department of Agriculture and Extension office.

Use this product only in accordance with the directions on the label. Timing and method of application, weather and crop conditions, mixtures with other materials, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all liability for any damage or loss of crop or other material not in strict accordance with directions given hereon.

E.P.A. Reg. No. 1202-247-AA