

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS  
DANGER**

Causes irreversible eye damage. May cause allergic skin reaction. Harmful if swallowed or inhaled. Do not get in eyes. Wear goggles or face shield and rubber gloves when handling. Avoid contact with skin and clothing. Avoid breathing of dust or vapor. Wash hands after use. Remove and launder clothing before use. Do not contaminate food or feedstuffs.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to water or wetlands. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift upon crops or animals. Do not contaminate water by cleaning of equipment or disposal of waste.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**REENTRY STATEMENT**

- Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.
- Do not enter treated areas without protective clothing until dusts have settled.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written or oral warnings must include the following information:

"**DANGER**" Area to be treated with captan on (date of application). Do not enter without appropriate protective clothing until dusts have settled. If accidentally exposed, consult the Statement of Practical Treatment portion of the pesticide label for first aid and medical attention recommendations."

**CROP RECOMMENDATIONS**

Apply when air is calm with an efficient dust applicator ensuring complete coverage with a light film of dust. Begin at first sign of infestation and repeat at 5-10 day intervals as needed to maintain control. Under severe disease conditions use the higher rate. For aerial or ground use.

**Agricultural Fruit and Nut Crop Uses**

- ALMOND** For brown rot, blight and blossom blight, etc. Apply 50-100 pounds per acre. Apply at blossom, bloom and post-bloom. Do not use within 12 days of harvest. When applied after post-bloom do not feed almonds hulls to dairy animals or animals being bred for slaughter.
- APPLES** (greenhouse) For bitter rot, brown rot, frog-eye leaf spot, black rot, scab, etc. Apply 40-75 pounds per acre. Apply in late cover and pre-harvest periods. For longleaf leaf spot use 40 to 60 pounds per acre in pre-harvest and blossom periods. Apply in first and second cover sprays and in 1 or 2 pre-harvest sprays as needed. For Braconid leaf spot, Resistant apply 22.75 pounds per acre. Apply in cover periods. For leaf spot - for primary infection apply 40-100 pounds per acre. Apply in pre-harvest, blossom, post-bloom and first cover periods. For secondary infection apply 20 to 30 pounds per acre. Apply in cover periods. For early blight apply 20-40 pounds per acre. Apply in cover periods. Note: Red Dickey and other sensitive varieties may be injured by early season applications.
- APRICOTS** (greenhouse) For brown rot, blight, etc. Apply 100 pounds per acre. Apply in late bloom, 75-90 percent petal fall and cover periods.
- BLACKBERRIES, RASPBERRIES** For anthracnose. Apply 40 pounds per acre. Apply in pre-harvest period when new canes are 8 to 8 inches long and in first cover period immediately after petal fall. For fruit rot apply 10 to 20 pounds per acre. Apply 3 to 5 days before harvest starts. Repeat at 10-day intervals and again 5 to 10 days later.
- BLUESBERRIES** For anthracnose. Apply 40-75 pounds per acre. Apply 20 pounds per acre. Begin at mid-bloom, repeat at 7 to 10-day intervals and re-apply.
- CANTALOUPES** (greenhouse) For anthracnose. Apply 100 pounds per acre. Apply at petal fall, bloom and in cover periods. For fruit rot apply 20 to 40 pounds per acre in 1st, 2nd, 3rd, 4th, 5th, 6th, 7th, 8th, 9th, 10th, 11th, 12th, 13th, 14th, 15th, 16th, 17th, 18th, 19th, 20th, 21st, 22nd, 23rd, 24th, 25th, 26th, 27th, 28th, 29th, 30th, 31st, 32nd, 33rd, 34th, 35th, 36th, 37th, 38th, 39th, 40th, 41st, 42nd, 43rd, 44th, 45th, 46th, 47th, 48th, 49th, 50th, 51st, 52nd, 53rd, 54th, 55th, 56th, 57th, 58th, 59th, 60th, 61st, 62nd, 63rd, 64th, 65th, 66th, 67th, 68th, 69th, 70th, 71st, 72nd, 73rd, 74th, 75th, 76th, 77th, 78th, 79th, 80th, 81st, 82nd, 83rd, 84th, 85th, 86th, 87th, 88th, 89th, 90th, 91st, 92nd, 93rd, 94th, 95th, 96th, 97th, 98th, 99th, 100th.
- CITRUS** (greenhouse) For anthracnose, etc. Apply 50 to 100 pounds per acre. Apply in bloom, blossom, post-bloom and cover periods. Repeat at 10-day intervals and again 5 to 10 days later.



**ACTIVE INGREDIENT**  
Captan (2,4,6-trichlorophenyl 4-cyanophenyl 1,2-dithiolane)  
**NET WT. 50 LBS.**

U.S. Patent Numbers 2,563,776 and 2,563,775  
E.P.A. Est. No.

**KEEP OUT OF REACH OF CHILDREN  
DANGER PELIGRO**

**PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que lo etiqueta le haya sido explicada ampliamente.**

**STATEMENT OF PRACTICAL TREATMENT**

- If in eyes—immediately flush eyes with plenty of water for at least 15 minutes. Call a physician.
- If swallowed—drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious person. Get medical attention.
- If inhaled—remove victim to fresh air. If not breathing give artificial respiration preferably mouth to mouth.
- If on skin—wash skin with soap and water.

**SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.**

- GRAPES** (house) For mold control on fruit on drying trays in field. Apply 20 pounds per acre by aerial application or 20 pounds per acre by ground equipment. Apply to fruit on trays. During damp weather make second application after trays are turned. For post-harvest application do not apply within 5 days of a pre-harvest application of captan.
- NECTARINES** (greenhouse) For brown rot, anthracnose, etc. Apply 100 pounds per acre. Apply in post-bloom, blossom, post-bloom and in late fall for anthracnose blight.
- PEACHES** (greenhouse) For brown rot, anthracnose, etc. Apply 40 to 100 pounds per acre. Apply in post-bloom, blossom, post-bloom and in late fall for anthracnose blight.
- PEARS** (greenhouse) For fruit rot, etc. Apply 40 to 60 pounds per acre. Apply in post-bloom, blossom, post-bloom and in late fall for anthracnose blight.
- PLUMS** (house) For brown rot, etc. Apply 40 to 100 pounds per acre. Apply at petal fall, 75-90 percent bloom, post-bloom and in cover sprays as necessary.
- QUINCE** For brown rot, etc. Apply 100 pounds per acre. Begin when disease pressure. Repeat at 7-10 day intervals. Do not apply within 7 days of harvest.
- RASPBERRIES** For anthracnose, etc. Apply 40 to 60 pounds per acre. Apply at bloom or when new canes are 8 to 10 inches long. Repeat 2 weeks later. Apply fall spray after canes are removed.
- BEANS** For anthracnose, etc. Apply 40 to 60 pounds per acre. Begin before disease starts or at first sign of disease. Repeat at 5 to 10-day intervals or as necessary to maintain control. For damping-off use 100 to 120 pounds per acre. Work into soil 3 to 4 inches of soil at planting time. Limit 50 pounds per acre use for one application. 60 pounds per acre as a preplant soil treatment.
- CANTALOUPES, CUCUMBERS, HONEYDEW, MELONS, PUMPKINS, SQUASH** (greenhouse and under) For anthracnose, etc. Apply 40 to 70 pounds per acre. Begin at first sign of disease. Repeat at 5 to 7 day intervals.
- CARROTS** For anthracnose, etc. Apply 40 to 60 pounds per acre. Begin when disease starts. Repeat at 5 to 10-day intervals.
- CELERY** For damping-off, etc. Apply 40 to 60 pounds per acre. Repeat as necessary. Do not exceed 40 pounds per acre. For use on plant bed treatment before transplanting apply 60-120 pounds per acre when all seeds have germinated. Do not exceed 120 pounds per acre.
- CORN** (house) For damping-off, etc. Apply 120 pounds per acre. Apply to soil before planting. Use one application 3 to 4 inches of soil. Do not apply within 7 days of harvest. Do not feed forage to dairy animals or animals being bred for slaughter.
- LETTUCE** For damping-off, etc. Apply 40 to 60 pounds per acre. Apply to soil before planting. Use one application 3 to 4 inches of soil. Do not apply within 7 days of harvest. Do not feed forage to dairy animals or animals being bred for slaughter.
- PEAS** For damping-off, etc. Apply 40 to 60 pounds per acre. Apply to soil before planting. Use one application 3 to 4 inches of soil. Do not apply within 7 days of harvest. Do not feed forage to dairy animals or animals being bred for slaughter.
- POTATOES** (Seed Potatoes) For brown rot, etc. Apply 100 pounds per acre. Apply 1 1/2 to 10 pounds per bushel. Dust seed pieces before planting, but pieces should be treated within 6 hours of cutting. 6 pounds in open crates before bagging. 10 pounds per bushel. Do not use treated seed for food or feed purposes.
- SPINACH** For damping-off, etc. Apply 40 to 120 pounds per acre. Pre-planting soil treatment. Work into upper 3 to 4 inches of soil. Or 50 to 70 pounds per acre in furrow over seed before covering. Limit 70 pounds per acre. 14 SEC. under test of soil as a furrow treatment. 120 pounds per acre as a broadcast preplanting soil treatment.
- TOMATOES** (Seed and greenhouse) For anthracnose, etc. Apply 40 to 60 pounds per acre. Apply to soil before planting. Use one application 3 to 4 inches of soil. Do not apply within 7 days of harvest. Do not feed forage to dairy animals or animals being bred for slaughter.

**STORAGE AND DISPOSAL**

- Do not contaminate water, food or feed by storage or disposal.
- Leaking of equipment should be stopped.
- When dumping is prohibited.
- STORAGE**
- Keep pesticide in original container.
- Store in cool, dry place.
- Protect from excessive heat.
- Do not store or transport near food or feed.
- Do not help with any spill leak fire or exposure involving material call day or night (800) 424-9300.
- PESTICIDE DISPOSAL**
- This product is acutely hazardous. Improper disposal of any pesticide spray mixture or residue is a violation of law. If these wastes cannot be disposed of by use accord to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.
- CONTAINER DISPOSAL**
- Completely empty bag into application equipment. Do not use container. Dispose of empty bag in a sanitary landfill. Do not burn, or if allowed by State and local authority, burning if burned stay out of smoke.

**PURCHASER CONDITIONS OF SALE AND WARRANTY**

PURECAP COMPANY warrants only that the product conforms to the other description on the label and is reasonably fit for the use stated on the label. Use in other circumstances with the directions under normal conditions of use. Critical and unimportant factors beyond PURECAP COMPANY'S control are beyond its control. Do not use in circumstances with the use of the product. Such results but do not result in damage to plants and crops to which the product is applied. Use of pesticide product cannot be PURECAP COMPANY and controlled by such as other factors or areas. Such use occurs even though the use is reasonable for the use stated herein and even though label directions followed. Therefore, the warranty does not extend to the use of the product contrary to the requirements of these directions or under conditions not stated herein. Do not use in circumstances with the use of the product. Such results but do not result in damage to plants and crops to which the product is applied. Use of pesticide product cannot be PURECAP COMPANY and controlled by such as other factors or areas. Such use occurs even though the use is reasonable for the use stated herein and even though label directions followed. Therefore, the warranty does not extend to the use of the product contrary to the requirements of these directions or under conditions not stated herein. Do not use in circumstances with the use of the product. Such results but do not result in damage to plants and crops to which the product is applied. Use of pesticide product cannot be PURECAP COMPANY and controlled by such as other factors or areas. Such use occurs even though the use is reasonable for the use stated herein and even though label directions followed.

