

DIRECTIONS FOR USE

Apply at the specified amount of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infestation and repeat at regular intervals when needed to maintain control, but observe use limits given on label.

CAUTION: Do not allow drift onto varieties of sorghum which are sensitive to phosphates, since injury may occur.

RECOMMENDATIONS

Field Crops:

ALFALFA: Apply 20 lbs. per acre to control alfalfa caterpillar, alfalfa webworm, Western Yellow-striped armyworm, beet armyworm, variegated cutworm, lygus bugs, stink bugs, tarnished plant bug. One application per cutting. Do not apply within 7 days of cutting or grazing.

COTTON: Apply 20-30 lbs. per acre to control cotton fleahopper, cotton leafworm, Western yellow-striped armyworm, beet armyworm, Southern garden leafhopper, black fleahopper complex, cotton leaf perforator, leaf roller, lygus bugs, stink bugs. Do not pasture treated fields within 14 days of application. Do not apply within 7 days of picking. Gavidox applications may occasionally cause marginal leaf burn.

SAFFLOWER: Apply 25-30 lbs. per acre to control armyworms, dipterous leaf miners, lygus bugs, variegated cutworm. Do not apply after flowering stage. Do not apply more than twice per year.


SUGAR BEETS: Apply 20-30 lbs. per acre to control beet webworm, variegated cutworm, dipterous leaf miners, alfalfa webworm, beet armyworm. Apply 30 lbs. per acre to control salt marsh caterpillar. Do not apply within 14 days of harvest. Tops harvested less than 28 days after treatment do not feed to livestock.

Vegetables:

BEANS (Dry and Snap): Apply 25-30 lbs. per acre to control Western bean cutworm, armyworms, dipterous leaf miners, lygus bugs, Mexican bean beetle, stink bugs, variegated cutworm. Do not apply within 14 days of harvest. Do not use treated vines as food for dairy or meat animals.

LIMA BEANS: Apply 20-30 lbs. per acre to control Western bean cutworm, armyworms, dipterous leaf miners, lygus bugs, Mexican bean beetle, stink bugs, variegated cutworm. Do not apply more than twice per season. Do not harvest for food, feed or forage within 14 days of treatment.

COWPEAS (Southern Peas, Black-eyed Peas, Crowder Peas): Apply 25-30 lbs. per acre to control armyworms, dipterous leaf miners, lygus bugs, stink bugs, variegated cutworm. Apply only twice per season. Do not apply within 14 days of harvest. Do not use vines as food for dairy or meat animals.



Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate	5%
INERT INGREDIENTS:	95%
	100%

*Dylox is reg. U.S. Pat. Off. by Farbenfabriken bayer, A.G. Chemagro Corporation Licensee

E.P.A. Reg. No. 1202-268-AA

CAUTION KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed, inhaled or absorbed through skin. Avoid getting on skin or in eyes. Avoid breathing dust. Always wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use. Do not contaminate feed or food.

If poisoning occurs, obtain prompt medical aid.
TO PHYSICIAN: Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Do not use on other crops used for food or forage. Use only according to label directions. Application at rates above those shown may result in illegal crop residues. Do not treat food crops grown in greenhouses.

Empty bags should be buried or burned. (Stay away from smoke). If the bags are burned, the ashes should be buried in a safe place.



BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER: Apply 20 lbs. per acre to control Western yellow-striped armyworm, imported cabbageworm, diamondback moth. Do not apply within 21 days of harvest.

CARROTS: Apply 20-30 lbs. per acre to control Western yellow-striped armyworm, variegated cutworm, dipterous leaf miners, beet armyworms, lygus bugs. Repeat as necessary. Do not use tops for food or feed. Do not apply within 28 days of harvest.

COLLARDS, LETTUCE: Apply 20 lbs. per acre to control variegated cutworm, armyworms, beet webworm, diamondback moth, dipterous leaf miners, salt-marsh caterpillar. Do not apply within 28 days of harvest.

PEPPERS: Apply 20-25 lbs. per acre to control dipterous leaf miners, pepper maggot, serpentine leaf miners. Repeat as necessary. Do not apply within 21 days of harvest.

PUMPKIN: Apply 25 lbs. per acre to control variegated cutworm, squash bug. Apply only once per season and not within 14 days of harvest.

TABLE BEETS: Apply 20-30 lbs. per acre to control variegated cutworm, alfalfa webworm, beet webworm, dipterous leaf miners, salt-marsh caterpillar, beet armyworm, lygus bugs. Repeat as necessary. Do not apply within 28 days of harvest. Do not eat tops.

TOMATOES: Apply 20-25 lbs. per acre to control serpentine leaf miners, tomato hornworms, dipterous leaf miners. Repeat as necessary. Do not apply within 21 days of harvest.

ENVIRONMENTAL CAUTIONS

This product is toxic to birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied where bees are actively visiting the area.

CONDITIONS OF SALE:

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.