

## DIRECTIONS FOR USE

**FRUIT AND NUTS CROPS:** Rates given in pints per 100 gallons unless specified. Aircraft—use equivalent rate in 10-50 gallons water per acre.

**APPLES:** 1 pint, Red-banded leaf roller. Do not apply after second cover spray or within 40 days of harvest.

**PEACHES:** 1.5 pints, plum curculio. Apply when plum curculio first appears and repeat every 10 to 14 days as necessary. Do not apply after second cover spray or within 40 days of harvest.

**PEACHES:** 2 pints, grasshopper. Do not apply after second cover spray or within 40 days of harvest.

**CITRUS** (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangeloes, Tangerines): 6 pints per acre, cutworms. Apply to ground under trees. 3 to 4 pints per 100 gals. water, grasshoppers. 1 pint per 100 gals. water, fire ants. Apply to trunk and main branches. Do not apply within 30 days of harvest. Do not feed treated forage to dairy animals or animals being finished for slaughter.

**CRANBERRIES:** 4 pints, armyworms. 3 pints, grasshoppers. Do not apply after fruit begins to form.

**PEARS:** 2 pints, grasshoppers. 1.2 pints, pear psylla, red banded leafroller. Do not apply within 40 days of harvest or after second cover spray.

**STRAWBERRIES:** 2-3 pints per acre, cutworm (climbing only) strawberry crown borer, strawberry weevil. Do not apply after fruit begins to form.

**FORAGE, FIELD, SEED AND VEGETABLE CROPS:** Rates given in pints per acre. Do not exceed recommended rates. Ground rig use 15-100 gallons water per acre. Aircraft 10-50 gallons water per acre.

**ALFALFA (Seed crops only):** 2½ to 3½ pints, alfalfa looper, alfalfa weevil, armyworm, green cloverworm. 3½ pints, climbing cutworm, grasshopper. 3 pints, alfalfa caterpillar, pea aphid, webworm. 2 pints, clover leaf weevil. 1½ pints spittlebug, alfalfa plant bug. Do not feed treated forage to dairy animals or animals being finished for slaughter.

**BARLEY, OATS, WHEAT:** 2 pints, armyworm, grasshopper. 3 pints, cutworm (climbing and surface feeding). Do not feed treated forage to dairy animals or animals being finished for slaughter. No restriction on the use of the grain.

**BEANS (Dry, Green, Lima):** 2 to 3 pints, grasshopper. 3 to 4 pints, garden webworm. 2 to 4 pints armyworm, mexican bean beetle, bean leaf beetle, cucumber beetle, blister beetle, flea beetle, cutworm (climbing and surface feeding), lygus bug, bean leaf skeletonizer, garden webworm, pea weevil. 3 pints, corn earworm, serpentine leafminer. 1.6 pints, thrips. 1 pint, bean leaf roller, bean leaf hopper. No limitation on use of shelled beans as human food. Do not apply sprays to green or snap beans after pods begin to form. Do not feed treated vines to dairy animals or animals being finished for slaughter.

**BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER:** 2 to 4 pints, cabbage looper, corn earworm, diamondback moth. 2 to 3 pints, flea beetle. 2 pints, green peach aphid, serpentine leaf miner, thrips. 1.6 pints, climbing cutworm, fall armyworm. Do not apply within 30 days of harvest or after edible parts start to form.

**CABBAGE:** 2 to 4 pints cabbage looper, corn earworm, diamondback moth. 2 to 3 pints, flea beetle. 2 pints, green peach aphid, serpentine leaf miner, thrips. 1.6 pints climbing cutworm, fall armyworm. Do not apply after heads start to form.

**CARROTS:** 2 to 4 pints, lygus bug. 2 to 3 pints, climbing cutworm, flea beetle. 2 pints, grasshopper. No time limitation.

**CELERY:** 1.6 pints cutworm (climbing and surface feeding), fall armyworm, green peach aphid, serpentine leaf miner, thrips. 4-5 pints, celery leaf tier. Do not apply after celery plants start to bunch or after plants are half mature.

**CORN:** 2 to 3 pints, chinch bug, cutworm (climbing), flea beetle, grasshopper. 2 to 2½ pints, webworm. Do not feed treated forage to dairy animals or animals being finished for slaughter. No limitation on use of grain.



## TOXAPHENE 8 LIQUID

### ACTIVE INGREDIENTS:

Toxaphene .....	72.5%
(Technical Chlorinated Camphene 67 - 69% Chlorine)	
NEEDLE INGREDIENTS .....	10.0%
	11.5%

8 Lbs. Toxaphene per gallon

E.P.A. Reg. No. 1202-271-AA

**WARNING**  
**KEEP OUT OF**  
**REACH OF CHILDREN**


Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies. Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.



LOS ANGELES, CALIFORNIA

NET CONTENT

GALS.





# TOXAPHENE 8 LIQUID

### ACTIVE INGREDIENTS

Toxaphene

Technical Concentrate Company, Inc., Dallas, Texas

### INERT INGREDIENTS

8.0% Total active per gallon

E.P.A. Reg. No. 120-071-AA

ACCEPTED  
AUG 14 1975

Do not use on any crop unless it is specifically listed on the label. Do not use on any crop unless it is specifically listed on the label. Do not use on any crop unless it is specifically listed on the label.



ANGELES, CALIFORNIA

**COTTON:** 2 to 4 pints/acre. Insects controlled: worm, cutworm, armyworm, bollworm, leafworm, beetle, grasshopper, yucca bug, caterpillar, flea hopper, thrips, 4 pints - link bug. Do not graze dairy animals or animals being finished for slaughter in fields treated later than 30 days after application. Do not use after boll opening.

**COWPEA:** (Including Field Peas) 1 pint/bean leafhopper, bean leafhopper, 2-3 pints cowpea caterpillar, yucca bug, 3 pints cutworms, armyworm, darkling ground beetle. No limitation on use of shelled peas as animal feed. Do not apply to cowpeas to be used as green snap after pod development. Do not feed treated peas to dairy animals or animals being finished for slaughter.

**EGGPLANT, PEPPERS:** 2 pints/acre. Insects controlled: potato beetle, cutworm, armyworm, 1 pint/eggplant leafhopper, leafhopper, serpentine leafhopper, 2 pints/cucumber beetle, 1 pint/eggplant flea beetle within 5 days of harvest.

**LETTUCE:** 3-5 pints/acre. Insects controlled: 4 pints/cucumber beetle, 3 pints/cutworm, armyworm. Do not apply after seedling stage. Do not use on lettuce heads to be marketed within 10 days of harvest.

**ONIONS:** 2 pints/acre. Do not apply to onion plants.

**PARSNIPS:** 1 pint/acre. Do not apply to parsnip plants.

**RICE:** 1 pint/acre. Insects controlled: armyworm, cutworm, leafhopper, grasshopper, stem borer, 1 pint/leafhopper, stem borer, 1 pint/leafhopper, stem borer, 1 pint/leafhopper, stem borer. Do not apply to rice plants until after the rice has been planted.

**RUTABAGUS:** 1 to 4 pints/acre. Insects controlled: cutworm, armyworm, 1 pint/eggplant leafhopper, 1 pint/eggplant flea beetle.

**SOYBEANS:** 1 pint/acre. Insects controlled: bean leafhopper, leafhopper, cutworm, armyworm, grasshopper, yucca bug, 1 pint/eggplant leafhopper, 1 pint/eggplant flea beetle, 1 pint/eggplant flea beetle, 1 pint/eggplant flea beetle, 1 pint/eggplant flea beetle. Do not apply to soybean plants until after the soybean has been planted.

**TOMATOES:** 1 to 4 pints/acre. Insects controlled: cutworm, armyworm, 1 pint/eggplant leafhopper, 1 pint/eggplant flea beetle, 1 pint/eggplant flea beetle, 1 pint/eggplant flea beetle, 1 pint/eggplant flea beetle. Do not apply to tomato plants until after the tomato has been planted.

**MAY BE FATAL IF SWALLOWED. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Keep away from children and domestic animals. Avoid contamination of feed and foodstuffs. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not feed forage crops treated with this material to dairy animals or to animals being finished for slaughter. Do not pasture other animals on forage within 3 weeks after treatment. Do not use on your plants. Do not use on cucurbits or prunes. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops rendered unfit for sale, use or consumption.**

### ENVIRONMENTAL CAUTIONS

Toxaphene is a highly toxic insecticide. It is highly persistent in the environment and is highly toxic to fish and aquatic life. It is also highly toxic to birds and mammals. It is highly toxic to bees and other beneficial insects. It is highly toxic to earthworms and other soil organisms. It is highly toxic to plants and animals. It is highly toxic to humans. It is highly toxic to the environment. It is highly toxic to all life forms.

### CONDITIONS OF SALE

This product is sold as a concentrate. It is to be diluted with water before use. It is to be used only on the crops and insects listed on the label. It is to be used only in accordance with the directions on the label. It is to be used only by persons who are properly trained and licensed to use pesticides. It is to be used only in accordance with the laws and regulations of the United States and the State of California. It is to be used only in accordance with the label instructions.