

DIRECTIONS FOR USE

NECTARINE: For the control of brown rot blossom blight (Monilinia) apply 60 lbs. per acre at pink bud and full bloom. For the control of brown rot fruit decay (Monilinia) apply at 60 lbs. per acre. Apply 18, 10 and 1 day before harvest. Do not apply later than one day before harvest.

PEACHES: For control of brown rot blossom blight (Monilinia) apply 60 lbs. per acre. Apply at 18, 10 and 1 day before harvest. For the control of rhizopus fruit rot apply 60 lbs. per acre. Apply one day before harvest. Do not apply later than one day before harvest.

SWEET CHERRY: For the control of brown rot blossom blight (Monilinia) apply 50 lbs. per acre. Apply at pop-corn, bloom, full bloom and petal fall. For the control of fruit decay (Monilinia) and rhizopus fruit rot apply 50 lbs. per acre. Apply 10 days and 1 day prior to harvest. Do not apply later than one day before harvest.

GRAPES: For the control of botrytis rot apply 30 lbs. per acre. Begin application while clusters are still loose and repeat application at two week intervals. Do not apply later than one day before harvest.

TOMATOES (Greenhouse): For the control of stem canker (botrytis) apply 30 lbs. per acre. Dust stem of plant from ground level up to a height of 18 to 24 inches. Repeat weekly or as necessary.

ONIONS, GARLIC (Not for use in North Central States): For control of white rot (sclerotium) apply 250-400 lbs. per acre. For direct seeding use mix into top 1.5 inches of soil one to two weeks before onions are seeded, for planting sets or buds apply broadcast and mix into top 1.5 inches, or apply same rates/43,560 linear feet of furrow to soil around sets or buds. Do not plant spinach as a follow-up crop on treated soil. When lettuce is to be planted as a follow-up crop, plow soil to 8 inches and cross disk the area prior to seeding.

LETTUCE: (Leaf, Greenhouse Only): For control of blight (botrytis rot) apply 30 lbs. per acre. Apply 7 days after transplanting. Repeat when plants are half mature. Do not apply to wilted plants or seedlings. Do not apply later than 14 days before harvest.



BOTRAN 6 DUST

ACTIVE INGREDIENTS:

2,6-Dichloro-4-nitroaniline6%

INERT INGREDIENTS94%
100%

Tuco Products Company — Division of the Upjohn Company
U.S. Pat. No. 3,119,736

EPA Reg. No. 1202-273-AA

CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid inhalation of dust. In case of contact, remove contaminated clothing and wash skin with soap and water.

Do not reuse container. Destroy when empty.



NET WEIGHT — 50 LBS.

ACCEPTED
11 SEP 1975
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO. 1202-273-AA

LETTUCE (Head, Field): For the control of botrytis rot and sclerotinia wilt, apply 33.3 pounds per acre. Apply at thinning, repeat seven to ten days later. Do not apply within 14 days of harvest.

ROSE, HYDRANGEA: For the control of botrytis gray mold, apply 30 lbs. per acre. Field—Apply BOTRAN 6 DUST prior to lifting for shipment or storage.

CHRYSANTHEMUM, GERANIUM: For the control of botrytis gray mold, apply 30 lbs. per acre. Greenhouse plants — Begin when disease is anticipated or first appears. Dust foliage and flowers at 7 to 14 day intervals. **Stock Cuttings:** Apply 8½ to 16½ lbs. as a uniform dust to cuttings and shipping cartons or cuttings in propagating bench, shipment or storage.

GLADIOLUS: For control of stromatinia dry rot apply 250-265 lbs. per 14,500 linear feet or row. Apply in a 12-inch band centered on furrow at time of planting. For control of botrytis blight apply 30 lbs. to bunched cut flower spikes, from one acre after harvesting and before storage. Use sufficient pressure to insure coverage between spikes.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

CONDITIONS OF SALE

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

LOT NO. _____ 4/75 CN

ACCEPTED WITH COMMENTS