

## DIRECTIONS FOR USE

For aircraft or ground rig use.

**BEANS** (Black-eyed peas, Cowpeas): Downy mildew, bean aphid, leaf miners, Mexican bean beetle, banded cucumber beetle — 42½ to 46½ pounds per acre. Begin at first sign of disease. Repeat 7- to 10-day intervals. Do not apply within 7 days of harvest.

**BEANS** (Snap or string): Anthracnose, downy mildew, rust, bean aphid, leaf miners, Mexican bean beetle, banded cucumber beetle — 26¼ to 46½ pounds per acre. Begin when plants are small. Repeat at 5- to 7-day intervals. Do not apply within 7 days of harvest.


**CABBAGE:** Alternaria leaf spot, downy mildew, Diamondback moth, aphids, imported cabbageworm — 26¼ to 40 pounds per acre. Begin 7- to 10-days after planting. Repeat at 3- to 5-day intervals in plant bed and field. Do not apply within 7 days of harvest.

**MELONS** (cantaloupes, Cassabas, Crenshaws, Honeydews, Honey Balls, Muskmelons, Persian Melons and Watermelons): Anthracnose, cercospora leaf spot, downy mildew, gummy stem blight, aphids, leafhoppers, mites, thrips, Serpentine leaf miners — 27½ to 46½ pounds per acre. Begin at late bunching, when vines begin to run, or when disease first appears. Repeat at 7- to 10-day intervals. Do not apply within 5 days of harvest.

**PEPPERS:** Anthracnose, cercospora leaf spot (frog-eye spot), ripe rot, aphids, leafhoppers, mites, thrips, Serpentine leaf miner — 53½ pounds as a dust/acre. Begin when disease threatens. Repeat at 7- to 10-day intervals. Do not apply within 5 days of harvest.

**POTATOES:** Early and late blights, aphid, flea beetle, Colorado potato beetle, Dipterous leaf miner, banded cucumber beetle, leafhoppers, Southern armyworms — 30 to 66½ pounds per acre. Begin when plants are 2 to 6 inches high. Repeat at 5- to 10-day intervals, more often under severe late blight conditions. Do not apply within 35 days of harvest.

**TOMATOES:** Cladosporium leaf mold, early and late blights, gray leaf spot (Stemphylium), septoria leaf spot, aphid, Dipterous leaf miner, banded cucumber beetle, Fall armyworms, Southern armyworms, vinegar fly — 30 to 53½ pounds per acre. Begin at first fruit cluster and repeat at 7- to 10-day intervals. Do not apply within 5 days of harvest.



## DIAZINON® MANEB DUST 3-4

### ACTIVE INGREDIENTS:

O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidyl) Phosphorothioate	.....3%
Maneb (Manganese Ethylenebisdithiocarbamate)	.....4%

INERT INGREDIENTS: .....93%

100%

DIAZINON® trademark of CIBA-GEIGY Corporation.

E.P.A. Reg. No. 1202-280-AA

## CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May be absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid storage near food and feed products.



NET WEIGHT — 50 LBS.

## ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Apply this product only as specified on this label.

Empty bags should be buried or burned. (Stay away from smoke). If the bags are burned, the ashes should be buried in a safe place.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

## CONDITIONS OF SALE:

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

LOT NO. \_\_\_\_\_ 7/75 CB