

DIRECTIONS FOR USE

Apply by ground rig or aircraft. A thorough application should be made. Do not exceed rate given for each crop.

APPLES, PEACHES, PEARS: Apply 25-150 lbs. per acre to control flea beetle, codling moth, gypsy moth, caterpillar, Fuller's rose beetle, plum curculio. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Remove excess residues by washing, brushing, or other effective means.

BEANS: (Dry Green, Lima, Snap) Apply 25-50 lbs. per acre to control Mexican bean beetle, bean leaf beetle, bean leaf roller, corn earworm, cucumber beetle. Remove excess by washing, brushing, or other effective means.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, COLLARD, MUSTARD, RADISHES, TURNIPS: Apply 25-100 lbs. per acre to control cabbage looper, climbing cutworm, corn earworm, flea beetle. Do not apply after edible parts start to form on broccoli, brussels sprouts, cabbage. Do not apply after seedling stage on cauliflower, collard, mustard greens, turnips. No time limitation on radishes.

CITRUS: (Grapefruit, Lemons, Limes, Oranges, Tangelos, Tangerines) Apply 25-100 lbs. per acre to control orange tortrix. No time limitation.

CANTALoupES, MELONS, WATERMELONS, CUCUMBER, SQUASH, PUMPKINS: Apply 25-100 lbs. per acre to control melon worms, pickleworm, cucumber beetle. Remove excess residues by washing, brushing or other effective means.



CRYOLITE-SULFUR 45-50 DUST

ACTIVE INGREDIENTS:

Cryolite 45
Sulfur 50

INERT INGREDIENT:

45
50
5
100

EPA Reg. No. 1292-253 AA

CAUTION KEEP OUT OF REACH OF CHILDREN

To protect fish and wildlife, do not contaminate lakes, ponds or streams with this material.

Empty bags should be buried or burned. Stay away from smoke. If the bags are burned, the ashes should be buried in a safe place.



NET WEIGHT

LBS

LOT NO

1-75 CN

CRANBERRIES: Apply 25-100 lbs. per acre to control cranberry weevil, gypsy moth, caterpillar, fruitworm, cranberry blossom worm, false armyworm. Remove excess residues by washing, brushing, or other effective means.

GRAPES: Apply 25-100 lbs. per acre to control leaf roller. Do not apply after fruit starts to form.

LETTUCE: Apply 15-100 lbs. per acre to control corn earworm, cabbage loopers. Do not apply after seedling stage.

PEPPERS: Apply 15-100 lbs. per acre to control pepper weevil. Remove excess residues by washing, brushing or other effective means.

STRAWBERRIES: Apply 25-100 lbs. per acre to control leaf roller. Remove excess residue by washing, brushing or other effective means.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

CONDITIONS OF SALE:

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.