

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

REENTRY STATEMENT

Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift.

Do not enter treated areas without protective clothing until dusts have settled.

Certain states may require more restrictive reentry intervals; consult your State Department of Agriculture for further information.

Written or oral warnings regarding use of protective clothing and accidental exposure must be given to workers who are expected to be in treated areas or in areas about to be treated.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

STORAGE

Keep pesticide in original container.

Store in cool, dry place.

Protect from excessive heat.

Do not store or transport near feed or food.

For help with any spill, leak, fire or exposure involving this material, call day or night (800) 424-9300.


PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at any approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

SUGGESTED USE

DIPEL 2 Dust is a biological insecticide preparation suitable for use in conventional dust application equipment as an aid in control of leaf chewing larvae (caterpillars) of certain lepidopterous species frequently found as pests on crop plants.

LOT NO. 8/84CD



ACTIVE INGREDIENTS:

Bacillus thuringiensis Berliner

Potency of 300 million

International Units per mg.

at least .48 Billion viable

spores per mg. 0.05%

INERT INGREDIENTS: 99.95%

100.0%

*Equivalent to potency of 0.14 Billion International Units per pound (quart) of this product.

DIPEL® is the trademark of the Agricultural and Veterinary Products Division, Abbott Laboratories.

EPA Reg. No. 1202-286-AA
E.P.A. Est. No.

KEEP OUT OF
REACH OF CHILDREN

CAUTION

WARNING: Reasonable care should be exercised to avoid inhalation or contact with open wounds.

ACCEPTED

JAN 3 1985

DISTRIBUTED
BY



WEST SACRAMENTO, CALIFORNIA

NET WEIGHT _____ LBS.

1202-286-AA 17

DIRECTIONS FOR USE

DIPEL 2 Dust is recommended for use by ground or by air. Apply when pests are first noticed and repeat as necessary. Thorough plant coverage is necessary for control.

Application Rate:

Broccoli, Cabbage, Cauliflower, Celery, Collards, Kale, Melons, Mustard Greens, Potatoes, Spinach, Tomatoes, Turnip Greens: For the control of cabbage looper and imported cabbageworm, apply 25-40 lbs. per acre.

COTTON:

For control of cabbage looper 29-40 lbs. per acre. For imported cabbage worm 25-40 lbs. per acre.

GRAPES:

For the control of grape leaf folder, apply 25-40 lbs. per acre.

LETTUCE:

For control of imported cabbage worm 29-40 lbs. per acre. For cabbage looper, 25-40 lbs. per acre.

PUREGRO CONDITIONS OF SALE AND WARRANTY

PureGro Company warrants that this product conforms to the chemical description on the label and is reasonably fit for purpose stated on such label. PureGro makes no other warranty, express or implied, of fitness or merchantability nor authorizes any agent or representative to make any variation or exception from this warranty.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of PureGro Company. Buyer assumes all risk of use, storage, or handling of this material not in strict accordance with directions given herewith.