

## DIRECTIONS FOR USE

**TOMATOES:** Apply 25 lbs. per acre to control drosophila, two-spotted mite, aphids, leaf-miner, tomato fruitworm, hornworm. Do not apply within 4 days of harvest.

**BEANS:** Lygus Bugs, Stink Bugs, Mexican Bean Beetle — 28 lbs. per acre. Apply thoroughly when insects appear and repeat every 7 to 10 days as necessary. Do not apply within 4 days of harvest.

**COTTON:** Salt-Marsh Caterpillar, Bollworms, Lygus, Cotton Fleahopper, Aphids, Cotton Leaf Perforator — 35 lbs. per acre. Apply as insects appear. Repeat at 7 to 10-day intervals while pests persist. Do not apply within 4 days of hand-picking.

**GRAPES:** Leafhoppers, Red-banded Leafroller — 25 lbs. per acre. Apply thoroughly when insects appear and repeat as necessary. Do not apply within 4 days of harvest.

**PEACHES:** Mites, Oriental Fruit Moth, Twig Borer — 50 lbs. per acre. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not apply within 4 days of harvest.

**PEACHES:** Mites, Oriental Fruit Moth, Twig Borer — 50 lbs. per acre. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not apply within 4 days of harvest.



## DIBROM® SEVIN® SULFUR 4-7.5-50 DUST

### ACTIVE INGREDIENTS

*Naled.....	4.0%
Carbaryl (1-naphthyl N-methylcarbamate) .....	7.5%
Sulfur .....	50.0%
INERT INGREDIENTS.....	38.5%
	100.0%

\*1,2-dibromo 2,2-dichloroethyl dimethyl phosphate trademark of Chevron Chemical Company, U. S. Patent No. 2,971,882. Trademark of Union Carbide Corporation, U. S. Patent No. 2,903,478

EPA Reg. No. 1202-291-AA / 202-29/

### CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or inhaled. May cause irritation of eyes, nose, and throat. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid storage near feed and food products.

Empty bags should be buried or burned. (Stay away from smoke). If the bags are burned, the ashes should be buried in a safe place.



LOS ANGELES, CALIFORNIA

NET WEIGHT

50

LBS.

LOT NO.

7/74 CN

## ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**CONDITIONS OF SALE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

### DIRECTIONS FOR USE

**TOMATOES:** Apply 25 lbs. per acre to control drosophila, two-spotted mite, aphids, leaf-miner, tomato fruitworm, hornworm. Do not apply within 4 days of harvest.

**BEANS:** Lygus Bugs, Stink Bugs, Mexican Bean Beetle — 28 lbs. per acre. Apply thoroughly when insects appear and repeat every 7 to 10 days as necessary. Do not apply within 4 days of harvest.

**COTTON:** Salt-Marsh Caterpillar, Bollworms, Lygus, Cotton Fleahopper, Aphids, Cotton Leaf Perforator — 35 lbs. per acre. Apply as insects appear. Repeat at 7 to 10-day intervals while pests persist. Do not apply within 4 days of hand-picking.

**GRAPES:** Leafhoppers, Red-banded Leafroller — 25 lbs. per acre. Apply thoroughly when insects appear and repeat as necessary. Do not apply within 4 days of harvest.

**PEACHES:** Mites, Oriental Fruit Moth, Twig Borer — 50 lbs. per acre. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not apply within 4 days of harvest.

**PEACHES:** Mites, Oriental Fruit Moth, Twig Borer — 50 lbs. per acre. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not apply within 4 days of harvest.



### DIBROM SEVIN SULFUR 4-7.5-50 DUST

#### ACTIVE INGREDIENTS

\*Naled 4.0%  
Carbaryl (1-naphthyl N-methylcarbamate) 7.5%  
Sulfur 50.0%

#### INERT INGREDIENTS

38.5%  
100.0%

\*1,2-dibromo-2,2-dichloroethyl dimethyl phosphate trademark of Chevron Chemical Company, U. S. Patent No. 2,971,882. Trademark of Union Carbide Corporation, U. S. Patent No. 2,903,478

EPA Reg. No. 1202-291-AA / 202-29/

### CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or inhaled. May cause irritation of eyes, nose, and throat. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid storage near feed and food products.

Empty bags should be buried or burned (Stay away from smoke). If the bags are burned, the ashes should be buried in a safe place.



NET WEIGHT 50

LBS LOT NO

774 CN

### ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**CONDITIONS OF SALE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

BEST DOCUMENT AVAILABLE