

**ACCEPTED**  
**JUN 9 1975**  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.



**MALATHION CRYOLITE  
 5-50 DUST**

**DIRECTIONS FOR USE**

**MELONS, CUCUMBERS, SQUASH:** Apply 30-45 lbs. per acre to control melon worms, pickleworms, cucumber beetle, aphid, spider mites. Do not apply within 1 day of harvest. Remove excess residues by washing, brushing or other effective means.

**BEANS:** (Dry, Green, Lima, Snap) Apply 30-35 lbs. per acre to control Mexican bean beetle, bean leaf beetle, bean leaf roller, corn earworm, cucumber beetle, leafhoppers, spider mites, Japanese beetle, aphids. Remove excess residues by washing, brushing, or other effective means. Do not apply within 1 day of harvest.

**BROCCOLI, CABBAGE, CAULIFLOWER:** Apply 30-60 lbs. per acre to control cabbage caterpillar, climbing cutworm, flea beetle, aphids, mites, imported cabbage worm, cabbage looper, diamondback moth. Do not apply after edible parts start to form on broccoli, cabbage. Do not apply after seedling stage on cauliflower.

**BRUSSELS SPROUTS:** Apply 30-50 lbs. per acre to control cabbage caterpillar, climbing cutworm, corn earworm, flea beetle, aphids, mites, imported cabbage worm, cabbage looper, diamondback moth. Do not apply after edible parts start to form on brussels sprouts.

**STRAWBERRIES:** Apply 25-40 lbs. per acre to control lygus bug, spittle bug, field crickets, thrips, potato leafhopper, strawberry leaf roller, white fly. Apply 30-40 lbs. per acre to control aphids, spider mites. Do not apply within 3 days of harvest. Remove excess residue by washing, brushing or other effective means.

**PUMPKINS:** Apply 25-40 lbs. per acre to control melon worms, pickleworm, cucumber beetle, aphid, spider mites. Do not apply within 3 days of harvest. Remove excess residues by washing, brushing, or other effective means.

**GRAPES:** Apply 30-55 lbs. per acre to control leaf roller, mealybugs, Japanese beetles, Drosophila, leafhopper. Do not apply after fruit starts to form.

**ACTIVE INGREDIENTS:**

Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate) .....5.0%

Cryolite (Sodium fluoaluminate) .....50.0%

**INERT INGREDIENTS:** .....45.0%

100.0%

E.P.A. Reg. No. 1202-297-AA

**CAUTION  
 KEEP OUT OF  
 REACH OF CHILDREN**

Avoid breathing of dust. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Harmful if swallowed.

**OBSERVE FOLLOWING USE PRECAUTIONS**

Observe use limitations given on the label for specific crops. Avoid dust drift to adjoining food and forage crops about to be harvested, or crops in bloom. Apply in early morning or evening when there is little or no wind. The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified.

Apply when air is calm using ground equipment or aircraft. Make thorough coverage of plants with a uniform deposit of dust. Apply at the first sign of infestation and repeat at regular intervals when needed to maintain control, but observe use limits given for specific crops.

Empty bags should be buried or burned (Stay away from smoke). If the bags are burned, the ashes should be buried in a safe place.



NET WEIGHT — 50 LBS

**LETTUCE:** Apply 30-50 lbs. per acre to control aphid, mites, cabbage looper, leafhoppers, corn earworm, alfalfa caterpillar, loopers, thrips. Do not apply after seedling stage.

**PEPPERS:** Apply 25-30 lbs. per acre to control aphids, pepper maggots, thrips, pepper weevil. Remove excess residues by washing, brushing or other effective means. Do not apply within 3 days of harvest.

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply to any area not specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**CONDITIONS OF SALE:**

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

LOT NO.

5/75 CN

