

DIRECTIONS FOR USE

Apply 20-40 lbs. per acre. Do not apply more than once per season. Do not apply within 14 days of harvest. Do not apply 1 day before harvest.

BEANS Mexican bean beetle, leafhoppers, bean leaf beetle, cucumber beetles, Western bean cutworm, flea beetles, Japanese beetle, velvet bean caterpillar, armyworms, tarnished plant bug, corn earworm. Apply 20-40 lbs. per acre. No time limitation.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI Flea beetle, harlequin bug, corn earworm, armyworms. For small loopers after heads form, use SEVIN every 5 to 7 days. Apply 20-40 lbs. per acre. Do not apply 3 days before harvest.

CORN Corn earworm. (Apply to silks as soon as they appear. Make 3 to 4 applications at 2-3 day intervals. Re-apply immediately after rain.) corn rootworm adults, European corn borer, fall armyworm, sap beetles, Japanese beetle, flea beetles, leafhoppers. Apply 20-40 lbs. per acre. No time limitation.

COTTON Thrips, flea beetle, leafhopper, leafworm, striped blister beetle, boll weevil, bollworm, fall armyworm, leaf perforator, leaf rollers, leafhoppers, tarnished plant bug, light to moderate numbers of Western lygus bugs (aphids will be repressed by repeated applications in a schedule), pink bollworm, salt marsh caterpillar, stink bugs, grasshoppers. Apply 30-40 lbs. per acre. No time limitations. Allow 7 days before grazing or harvesting crop residue for feed or bedding for dairy or meat animals.

LEAFY VEGETABLES AND ROOT CROPS (Chinese Cabbage, Collards, Dandelion, Endive, Kale, Leaf Lettuce, Mustard Greens, Parsley, Spinach, Swiss Chard). Apply 20-40 lbs. per acre to control flea beetles, leafhoppers, harlequin bug, six-spotted leafhopper, aster yellows vector, corn earworm, armyworms, lygus and stink bugs. Do not apply within 14 days of harvest.

PEAS Leafhoppers, Colorado potato beetle, armyworms, grasshoppers. Apply 20-30 lbs. per acre. No time limitation.

POTATO Colorado potato beetle, flea beetles, leafhoppers. Apply 20-40 lbs. per acre. Foliage application only.

SUGAR BEETS Leafhopper, armyworm, webworm, grasshoppers. Apply 20-40 lbs. per acre. Do not apply 14 days before harvest.

EDIBLE PLANT PEPPER Tomato fruitworm, fall armyworm, leaf worm, European corn borer, lace bugs, Colorado potato beetle, flea beetles, leafhopper. Apply 20 lbs. per acre. No time limitation.

CUCUMBERS, SQUASH, PUMPKIN, MELONS, PEPPERS Leafhopper, cucumber beetle, squash bug, flea beetle, leafhopper. Apply 20 lbs. per acre. No time limitation.

HEAD LETTUCE, HORSE RADISH, PARSNIP, RUTABAGA, RUTABAGAS Apply 20-40 lbs. per acre to control flea beetles, leafhoppers, harlequin bugs, six-spotted leafhopper, leaf yellow vector, corn earworm, armyworm, lygus and stink bug. Do not apply within 3 days of harvest.

BEETS Apply to TURNIPS. Apply 20 lbs. per acre to control flea beetles, leafhoppers, harlequin bug, six-spotted leafhopper, leaf yellow vector, corn earworm, armyworm, lygus and stink bug. Do not apply within 3 days of harvest.

Apply 20-40 lbs. per acre. Do not apply more than once per season. Do not apply within 14 days of harvest. Do not apply 1 day before harvest.



SEVIN[®] 5 DUST

ACTIVE INGREDIENT

Carbaryl (1-naphthyl N-Methyl carbamate)

5%

INERT INGREDIENTS

95%

100%

EPA REG. NO. 1202-300-AA

Registered trademark of Chemagro Carbide Corporation for the active ingredient (Carbaryl).

CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after using. Avoid storage near feed and food products.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

EMPTY BAGS SHOULD BE BURIED OR BURNED. STAY AWAY FROM SMOKE IF BUGS ARE BURNED IN THE ASHES. DO NOT BE BURIED IN A TREE HOLE.



DAVID A. F. R. A.

Apply 20-40 lbs. per acre. Do not apply more than once per season. Do not apply within 14 days of harvest. Do not apply 1 day before harvest.

APPLES, PEARS Codling moth, green apple aphid, pear psylla, orange tortrix, California pear slug. Apply 70-120 lbs. per acre. Do not apply 1 day before harvest.

CHERRY, PLUM, PRUNE Cherry fruit fly, fruit tree leaf roller, peach twig borer, orange tortrix, Western tussock moth. Apply 80-120 lbs. per acre. Do not apply 1 day before harvest.

GRAPES Grape leaf folder, grape leafhopper, grape leaf skeletonizer, grape berry moth, Japanese beetle, June beetles, red-banded leaf roller. Apply 40 lbs. per acre. No time limitation.

PEACHES, NECTARINES Oriental fruit moth, peach twig borer, codling moth, orange tortrix, tussock moth, fruit tree leaf roller, Japanese beetle. Apply 80-120 lbs. per acre. Do not apply 1 day before harvest for peaches and 3 days before harvest for nectarines.

Storage and Feed Crops

ALFALFA, CLOVERS, PEANUTS, SOYBEANS, COWPEAS Blister beetles, Mexican bean beetle, thrips, leafhoppers, three-cornered alfalfa hopper, cucumber beetles, bean leaf beetle, Japanese beetle, alfalfa caterpillar, velvet bean caterpillar and green cloverworm, armyworms, webworms, corn earworm, grasshoppers. Apply 20-30 lbs. per acre. No time limitation.

FORAGE GRASSES - PRAIRIE Apply 5-30 lbs. per acre to control armyworms, thrips, flea beetles, grasshoppers, grasshoppers, grasshoppers. Do not apply 14 days before harvest.

30 Broomcorn, Grain and Forage Armyworms, webworms, corn earworm, sugarcane borer, Colorado potato beetle. Apply 30 lbs. per acre 5 to 7 days after heads have emerged from field. Grasshoppers. Apply 30-40 lbs. per acre. Do not apply 21 days before harvest for grain. No time limitation for forage.

Apply 20-40 lbs. per acre. Do not apply more than once per season. Do not apply within 14 days of harvest. Do not apply 1 day before harvest.